



**QUEEN'S
UNIVERSITY
BELFAST**

The biology of habitat dominance; can microbes behave as weeds?

Cray, J. A., Bell, A. N. W., Bhaganna, P., Mswaka, A. Y., Timson, D. J., & Hallsworth, J. E. (2013). The biology of habitat dominance; can microbes behave as weeds? *Microbial Biotechnology*, 6(5), 453-492.
<https://doi.org/10.1111/1751-7915.12027>

Published in:
Microbial Biotechnology

Document Version:
Publisher's PDF, also known as Version of record

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

Copyright 2013 the authors.

This is an open access article published under a Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Minireview

The biology of habitat dominance; can microbes behave as weeds?

Jonathan A. Cray, Andrew N. W. Bell, Prashanth Bhaganna, Allen Y. Mswaka, David J. Timson and John E. Hallsworth*

School of Biological Sciences, MBC, Queen's University Belfast, Belfast, BT9 7BL, Northern Ireland, UK.

Summary

Competition between microbial species is a product of, yet can lead to a reduction in, the microbial diversity of specific habitats. Microbial habitats can resemble ecological battlefields where microbial cells struggle to dominate and/or annihilate each other and we explore the hypothesis that (like plant weeds) some microbes are genetically hard-wired to behave in a vigorous and ecologically aggressive manner. These 'microbial weeds' are able to dominate the communities that develop in fertile but uncolonized – or at least partially vacant – habitats via traits enabling them to out-grow competitors; robust tolerances to habitat-relevant stress parameters and highly efficient energy-generation systems; avoidance of or resistance to viral infection, predation and grazers; potent antimicrobial systems; and exceptional abilities to sequester and store resources. In addition, those associated with nutritionally complex habitats are extraordinarily versatile in their utilization of diverse substrates. Weed species typically

deploy multiple types of antimicrobial including toxins; volatile organic compounds that act as either hydrophobic or highly chaotropic stressors; biosurfactants; organic acids; and moderately chaotropic solutes that are produced in bulk quantities (e.g. acetone, ethanol). Whereas ability to dominate communities is habitat-specific we suggest that some microbial species are archetypal weeds including generalists such as: *Pichia anomala*, *Acinetobacter* spp. and *Pseudomonas putida*; specialists such as *Dunaliella salina*, *Saccharomyces cerevisiae*, *Lactobacillus* spp. and other lactic acid bacteria; freshwater autotrophs *Gonyostomum semen* and *Microcystis aeruginosa*; obligate anaerobes such as *Clostridium acetobutylicum*; facultative pathogens such as *Rhodotorula mucilaginosa*, *Pantoea ananatis* and *Pseudomonas aeruginosa*; and other extremotolerant and extremophilic microbes such as *Aspergillus* spp., *Salinibacter ruber* and *Haloquadratum walsbyi*. Some microbes, such as *Escherichia coli*, *Mycobacterium smegmatis* and *Pseudoxylaria* spp., exhibit characteristics of both weed and non-weed species. We propose that the concept of nonweeds represents a 'dustbin' group that includes species such as *Synodopsis* spp., *Polypaecilum pisce*, *Metschnikowia orientalis*, *Salmonella* spp., and *Caulobacter crescentus*. We show that microbial weeds are conceptually distinct from plant weeds, microbial copiotrophs, *r*-strategists, and other ecophysiological groups of microorganism. Microbial weed species are unlikely to emerge from stationary-phase or other types of closed communities; it is open habitats that select for weed phenotypes. Specific characteristics that are common to diverse types of open habitat are identified, and implications of weed biology and open-habitat ecology are discussed in the context of further studies needed in the fields of environmental and applied microbiology.

Received 11 October, 2012; accepted 3 December, 2012. *For correspondence. E-mail j.hallsworth@qub.ac.uk; Tel. +44 (0)28 9097 2314 (direct line); Fax +44 (0)28 9097 5877.

Microbial Biotechnology (2013) 6(5), 453–492

doi:10.1111/1751-7915.12027

Funding Information Funding was received from the Department of Agriculture and Rural Development (Northern Ireland), the research programme of the Kluyver Centre for Genomics of Industrial Fermentation which is part of the Netherlands Genomics Initiative/Netherlands Organization for Scientific Research, Biotechnology and Biological Sciences Research Council (BBSRC) for a Daphne Jackson Fellowship, BBSRC and Natural Environment Research Council United Kingdom projects BBF0034711 and NE/E016804/1 (respectively), and the Beaufort Marine Research Award for Marine Biodiscovery which is carried out under the Sea Change Strategy and the Strategy for Science Technology and Innovation (2006–2013), with the support of the Marine Institute, Ireland, and J.E.H. received a Prof. B. D. Tilak Fellowship from Institute of Chemical Technology-Mumbai, India.

Introduction

As the collective metabolism and ecological activities of microorganisms determine the health and sustainability of life on Earth it is essential to understand the function of

both individual microbes and in their communities. The cellular systems of an insignificant portion of microbial species have been intensively characterized at the levels of biochemistry, cell biology, genomics and systems biology; and a substantial body of (top-down) studies has been published in the field of microbial ecology in relation to species interactions, community succession and environmental metagenomics. There are, however, some fundamental questions that remain unanswered in relation to the behaviour of microbial species within communities: what type of biology, for instance, enables microbes to dominate entire communities and their habitats and thereby determine levels of biodiversity in specific environments?

Plant species that primarily inhabit freshly disturbed habitats – known as weeds – are characterized by vigorous growth; tolerance to multiple stresses; exceptional reproduction, dispersal and survival mechanisms; a lack of specific environmental requirements; production of phytotoxic chemicals; and/or other competitive strategies (Table 1). The biology of plant weeds, including their phenotypic and genetic traits, has been the subject of extensive study over the past 100 years and is now relatively well characterized (Table 1; Long, 1932; Anderson, 1952; Salisbury, 1961; Hill, 1977; Mack *et al.*, 2000; Schnitzler and Bailey, 2008; Bakker *et al.*, 2009; Chou, 2010; Wu *et al.*, 2011; Liberman *et al.*, 2012). We hypothesize that weed biology is also pertinent to microbial species, but that the defining characteristic of microbial weeds is their ability to *dominate* their respective communities. In the field of plant ecology it is open habitats (such as freshly exposed fertile soil) that facilitate the emergence of weed species both within specific ecosystems and across evolutionary timescales. We propose that weed behaviour is equally prevalent in some microbial habitats, that open microbial habitats promote the emergence of microbial weed species, and that microbial weed biology represents a potent ecological and evolutionary mechanism of change for some microbial species and their communities.

This article focuses on the following questions: (i) what types of substrate or environment facilitate the emergence of dominant microbial species, (ii) are there archetypal weed species that can consistently dominate microbial communities, (iii) what are the stress biology, nutritional strategies, energy-generating capabilities, antimicrobial activities and other competitive strategies of microbial weeds, (iv) which components and characteristics of microbial cells form the mechanics of weed biology, (v) what are the properties of open (microbial) habitats that promote the emergence of weed species, and (vi) how are microbial weeds conceptually distinct from plant weeds, and other ecophysiological groups of microorganism?

Dominance within microbial communities

The microbial diversity of specific habitats, as well as that of Earth's entire biosphere, is an area of intensive scientific interest that has implications within the fundamental sciences and for drug discovery, biotechnology and environmental sustainability. Biodiversity provides the material basis for competition between microbial taxa, species succession in communities, and other aspects of microbial ecology that collectively impact the evolution of microbial species (Cordero *et al.*, 2012). We believe that certain microbes are genetically, and thus, metabolically hardwired to dominate communities (monopolizing available resources and space) and thereby reduce biodiversity within the habitat. Microbial communities can reach a stationary-phase or climax condition with a relatively balanced species composition (McArthur, 2006). However whether tens or thousands of microbial species are initially present (Newton *et al.*, 2006) the community can become dominated by a single – or two to three – species (Fig. 1; Table S1; Randazzo *et al.*, 2010; Cabrol *et al.*, 2012). For example the indigenous microflora of olives is comprised of significant quantities of phylogenetically diverse species such as *Aureobasidium pullulans*, *Candida* spp., *Debaryomyces hansenii*, *Metschnikowia pulcherrima*, *Pichia guilliermondii*, *Pichia manshurica*, *Rhodotorula mucilaginosa* and *Zygowillipsis californica* (Fig. 1Bi; Nisiotou *et al.*, 2010). Nisiotou and colleagues (2010) found that species succession and the stationary-phase community during olive fermentations are determined by one dominating genus, *Pichia*, three species of which account for almost 100% of the community by 35 days (see Fig. 1Biii). The most prevalent of these, *Pichia anomala*, is a microbial generalist that is not only able to inhabit ecologically and nutritionally distinct environments (e.g. palm sugar, cereals, silage, oil-contaminated soils, insects, skin, various marine habitats; see Walker, 2011) but can also dominate the communities that develop in diverse habitat types (Table S1; Tamang, 2010; Walker, 2011). Studies of Pecorino Crotonese cheese fermentations reveal that, even though the curd contains > 300 bacterial strains, the community is eventually dominated by *Lactobacillus rhamnosus* (Randazzo *et al.*, 2010).

Species that are able to dominate communities that develop in open habitats are not restricted to copiotrophs, generalists or any established ecophysiological grouping of microorganism (Table S1). Furthermore dominating species span the Kingdoms of life and, collectively, can be observed in most segments of the biosphere; from bio-aerosols to solar salterns, oceans and freshwater to sediments and soils, the phyllosphere and rhizosphere, to those with molar concentrations of salts or sugars (see Table S1). The phenomenon can also be observed during food-production, food-spoilage and waste-decomposition

Table 1. A comparison of plant weeds and microbial weeds.^a

	Plant weeds	Microbial weeds	References (plants/ microbes)
<i>Definition</i>	Plant species that primarily grow in open (plant) habitats ^b	Microbial species able to dominate communities that develop in open (microbial) habitats ^c	Baker, 1965; Hill, 1977/ current article; Cordero <i>et al.</i> , 2012
<i>Ability to rapidly occupy available space</i>	Important trait	Essential trait	Hill, 1977/current article
<i>Overall vigour^d and competitive ability</i>	Typically vigorous; competitive ability usually superior to that of crop plants	Exceptional vigour and competitive ability	Hill, 1977/Figs 1 and 2; Tables S1 and 2–5
<i>Primary competitors</i>	Intra-specific competition can be stronger than inter-specific competition	Intra- and inter-specific competition are likely to be stronger if dominance results from efficient resource acquisition or antimicrobial substances respectively	Hill, 1977/current article
<i>Resource acquisition and storage</i>	Acquisition and/or storage of water, light, nutrients etc. typically excellent	Acquisition and/or storage of nutrients, water and/or light typically excellent	Hill, 1977; Oatham, 1997; Sadeghi <i>et al.</i> , 2007/ Table 5
<i>Reproduction and dispersal</i>	Exceptional ^e	May not differ from non-weed species	Hill, 1977/current article
<i>Obligatory associations or interactions with other organisms</i>	Uncommon/atypical	Uncommon/atypical	Hill, 1977/current article
<i>Ubiquity</i>	Many species are environmentally ubiquitous	Not necessarily widespread outside the habitat which they dominate	Hill, 1977/current article
<i>Germination</i>	No special requirements; high tolerance to diverse stresses (see below)	No special requirements; high tolerance to diverse stresses (see below)	Hill, 1977/current article
<i>Longevity of dormant propagules</i>	Exceptional	May not differ from non-weeds	Hill, 1977/current article
<i>Able to regenerate from fragments</i>	Yes, especially root fragments	Not relevant	Hill, 1977/current article
<i>Maintenance of growth rate during nutrient limitation</i>	Some species, e.g. black-grass (<i>Alopecurus mysuroides</i>) on low-potassium soils	Many (photosynthetic as well as some heterotrophic) species	Hill, 1977/Table 5; Benning, 1998
<i>Capable of creating a zone-of-inhibition</i>	Not known	Yes, in some instances	[Not known]/Fleming, 1929
<i>Secretion of toxic substances</i>	Some species produce phytotoxic inhibitors (exuded from leaves, roots, and/or necromass)	All species produce antimicrobial toxins (secreted into the extracellular environment) ^f	Chou, 2010/Table 6
<i>Production of volatile organic compounds (VOCs) that act as stressors</i>	Some hydrophobic and highly chaotropic compounds act as phytotoxic stressors (e.g. β -caryophellene)	Many species produce a spectrum of VOCs that can inhibit metabolic activity at concentrations in the nanomolar or low millimolar range ^f	Wang <i>et al.</i> , 2010; Inderjit <i>et al.</i> , 2011; Tsubo <i>et al.</i> , 2012/Table 6
<i>Secretion of biosurfactants that act as stressors</i>	Not known	Some species produce biosurfactants; substances that can solubilize hydrophobic substrates and act as cellular stressors ^f	[Not known]/Tables 3 and 5; Nitschke <i>et al.</i> , 2011
<i>Bulk production, and secretion, of moderately chaotropic substances</i>	Not known	Some species produce ethanol, butanol, acetone etc. that are potent cellular stressors at millimolar and molar concentrations ^f	[Not known]/Table 6
<i>Saprotrophic activity resulting in toxic/stressful breakdown products</i>	Not known	Some species degrade macromolecules to release inhibitory catabolic products (e.g. chaotropic stressors such as phenol, catechol and vanillin ^g)	[Not known]/Park <i>et al.</i> , 2004; 2007; Zuroff and Curtis, 2012
<i>Acidification of habitat</i>	Not known	Some species, via bulk production of organic acids ^f	[Not known]/Table 6
<i>Secretion of aggressive enzymes</i>	Unlikely	Some species ^f	[Not known]/Table 6; Walker, 2011
<i>Tolerance to environmental stresses^h</i>	Tolerant to multiple, habitat- relevant stress parameters	Tolerant to multiple, habitat-relevant stress parameters	Hill, 1977/Fig. 2, Tables 2–4 and 6
<i>Resistance to toxins and/or stressors of biotic origin^h</i>	Likely	Yes, most species	Hill, 1977/Fig. 2; Tables 2–4 and 6

Table 1. *cont.*

	Plant weeds	Microbial weeds	References (plants/ microbes)
<i>Phenotypic plasticity and intra-specific variation</i>	Greater than that of crop plants	Likely to be above average (e.g. in relation to stress biology)	Hill, 1977/current article; see references in Todd <i>et al.</i> , 2004
<i>Resistance to viruses, predators, and/or grazers</i>	Some species can tolerate, resist, and/or repel viruses, sucking insects and grazing animals; and may contain poisonous substances – or have morphological adaptations such as spines – that minimize losses through grazing	Some strains/species can tolerate, resist, repel and/or avoid viruses, grazing plankton and arthropods etc; may contain toxic metabolites or have behavioural adaptations – e.g. formation of large microcolonies – that minimize losses through grazing/predation; and may inhabit environments that are too extreme for predatory species	Thurston <i>et al.</i> , 2001; Ralphs, 2002; Shukurov <i>et al.</i> , 2012/ Table 6
<i>Karyotype</i>	Commonly of hybrid origin, polyploid or aneuploid	Frequency of hybrid origin, polyploidy and aneuploidy not yet tested; may contain plasmids that contribute to weediness	Anderson, 1952/Tables 3 and 5
<i>Proportion of total species</i>	Minute (several hundred species)	Not yet established (likely to be minute)	Hill, 1977/current article
<i>Association with human activity</i>	Evolution and ecology have been strongly favoured by land disturbance that creates and open (plant) habitats ⁱ	Evolution is likely to have occurred primarily in nature; but habitat dominance is commonplace in both natural and manmade environments	Anderson, 1952; Hill, 1977/Fig. 2; Tables S1 and 7
<i>Problematic for human endeavours</i>	Yes, due to loss of crop-plant yields ^j ; toxicity/harm to livestock; role as hosts for crop pests and diseases, etc.	Yes, due to spoilage of foods, drinks; some species are pathogens of crop plants, livestock or humans; may emerge to take over industrial fermentations and other processes ^k	Hill, 1977/Table S1
<i>Value to humankind</i>	Disproportionately important (origin of many crop-plant species; plant weeds have food value for humans or livestock; model organisms for research)	Disproportionately important (key sources of bulk chemicals and antimicrobials ^l ; for food and drinks production, and other biotechnological applications)	Hill, 1977; Koornneef and Meinke, 2010/ Table S1; see <i>Concluding remarks</i>

a. Typical traits.

b. Plant weeds may not be the dominating plant species.

c. Not necessarily the primary habitat of the microbial weed species.

d. Vigour can result from a combination of traits such as fast growth, stress tolerance, and immunity to toxins and diseases (see Tables 2–6).

e. Plant weeds typically have a high output of seeds under both favourable and poor growth-conditions, and can produce seed early (prior to maturity of the parent plant; Hill, 1977).

f. Each microbial weed species may secrete diverse antimicrobial substances which can be the primary factor in habitat dominance (see Table 7; Hallsworth, 1998; Walker, 2011). For examples of the concentrations at which moderately chaotropic stressors, potent chaotropes and hydrophobic stressors can inhibit cellular systems see Bhaganna and colleagues (2010). Toxins such as antibiotics, and VOCs, biosurfactants, and other stressors can have additional key roles in the cellular biology and ecology of microbes.

g. See Hallsworth and colleagues (2003a); Bhaganna and colleagues (2010).

h. Some stressors of biotic origin are structurally identical to and/or exert the same stress mechanisms as stressors of abiotic origin (e.g. see Fig. 2; Tables 2–4 and 6).

i. Agricultural practices and crop-plant life-cycle can also select for early seed production and other weed traits (Hill, 1977).

j. Crop-yield losses caused by plant weeds are equivalent to those caused by plant pests or plant diseases (Cramer, 1967).

k. Open (microbial) habitats are not only the starting point for food and drinks fermentations, but many foodstuffs themselves represent open habitats and it is frequently microbial weed species that cause spoilage and therefore determine shelf-life. Open habitats located on and within host organisms can be readily invaded by microbial weed species including those with pathogenic activities (Cerdeño-Tárraga, *et al.*, 2005). This has given rise to the use of probiotics (Molly *et al.*, 1996; Bron *et al.*, 2012) and a practice of preserving the skin microflora in babies and infants by avoiding use of soaps or detergents (Capone *et al.*, 2011).

l. Antimicrobial substances are used for various applications; as biofuels, biocides, pharmaceuticals, flavour compounds, food preservatives, etc.

processes where resource-rich habitats are available for colonization (see Table S1). The materials present in and the microbes living on these substrates are ultimately derived from the natural environment so community successions resulting in single-species dominance are likely to be commonplace in comparable open habitats within

natural ecosystems (Table S1; Senthilkumar *et al.*, 1993; Nisiotou *et al.*, 2007; Reynisson *et al.*, 2009; Rajala *et al.*, 2011). Microbial weed species are able to dominate open habitats that progress to a closed condition (e.g. fermenting fruit juice) as well as those that remain in a more or less perpetually open state (e.g. the surface-fluid film of




Fig. 1. Percentage species composition during olive fermentations at (i) 2 days, (ii) 17 days and (iii) 35 days over a range of conditions: (A) with added NaCl (6% w/v); (B) added NaCl and glucose (6% and 0.5% w/v respectively); (C) added NaCl and lactic acid (6% w/v and 0.2% v/v respectively); and (D) NaCl, glucose and lactic acid (6% and 0.5% w/v, and 0.2% v/v respectively). Data (from Nisiotou *et al.*, 2010) were obtained by DGGE.

sphagnum moss and rhizosphere of plants growing in moist soils); see Table S1. Weeds emerge in habitats where competing cells are in close proximity (such as biofilms; Table S1; Rao *et al.*, 2005) as well as those where there is limited mechanical contact between cells, such as the aqueous habitats of planktonic species (Table S1; Collado-Fabbri *et al.*, 2011; Trigal *et al.*, 2011; Leão *et al.*, 2012); and in both extreme and non-extreme environments (Table S1). Some microbes emerge as a/the dominating species in diverse types of habitat including specialists such as *Lactobacillus* (that do so, in part, by acidifying their environment) and generalists such as *Aspergillus*, *Pichia* and *Pseudomonas* species (Table S1; Steinkraus, 1983; Tamang, 2010). For weed species such as *Saccharomyces cerevisiae* (high-sugar environments), *Haloquadratum walsbyi* (aerobic, hyper-saline brines), and *Gonyostomum semen* and *Microcystis*

aeruginosa (eutrophic freshwater) ability to dominate can be restricted to a specific type of habitat. Most of these microbes do not populate their respective habitat as a one-off or chance event; they can consistently dominate their microbial communities and do so regardless of their initial cell number (Table S1; for *P. anomala* see Fig. 1; for *S. cerevisiae* see Pretorius, 2000; Renouf *et al.*, 2005; Raspor *et al.*, 2006; for *L. rhamnosus* see Randazzo *et al.*, 2010). It is therefore intriguing to consider what combination(s) of traits can give rise to a weed phenotype.

Cellular biology of microbial weeds

Stress tolerance and energy generation

During habitat colonization, even in apparently stable, homogenous environments such as sugar-based

solutions, there are dynamic/drastring changes in stressor concentrations, antimicrobials and stress parameters (D'Amore *et al.*, 1990; Hallsworth, 1998; Eshkol *et al.*, 2009; Fialho *et al.*, 2011; see also below). As a result of these fluctuations microbes are exposed to multiple stresses some of which present potentially lethal challenges to the cell. In soil and sediments, for example the water activity can oscillate from low to high and high to low; these substrates typically cycle between desiccation and rehydration and are subjected to changes in solar radiation, temperature, and even freezing and thawing. Soils and sediments can be physically heterogeneous such that water availability, solute concentrations and other stress parameters exhibit profound spatial fluctuations. In addition, intracellular metabolites and extracellular substances (of both biotic and abiotic origin) impose stresses due to ionic, osmotic, chaotropic, hydrophobic and other activities of solutes (see Brown, 1990; Hallsworth *et al.*, 2003a; 2007; Lo Nostro *et al.*, 2005; Bhaganna *et al.*, 2010; Chin *et al.*, 2010). On the one hand resilience to stress undoubtedly contributes to competitive ability but on the other a vigorous growth phenotype and reinforced stress biology also require extraordinary levels of energy generation. Furthermore, microbes that can achieve a greater net gain in energy than competing species will have an ecological advantage (Pfeiffer *et al.*, 2001). Here we suggest that species able to dominate microbial communities are exceptionally well equipped to resist the effects of multiple stress parameters, and that many have high-efficiency energy-generation systems.

We tested the hypothesis that microbial weeds are exceptionally tolerant to habitat-relevant stresses for two species; the soil bacterium *Pseudomonas putida* (Table 2) and the specialist yeast *S. cerevisiae* (see below). Environments in which *P. putida* is a major ecological player (Table S1) typically contain an array of chemically diverse aromatics and hydrocarbons that induce chaotropicity-mediated stresses (Hallsworth *et al.*, 2003a; Bhaganna *et al.*, 2010; Cray *et al.*, 2013). This bacterium is not only renowned for its metabolic versatility, but is also known for its tolerance to specific solutes and solvents (Hallsworth *et al.*, 2003a; Domínguez-Cuevas *et al.*, 2006; Bhaganna *et al.*, 2010). We compared the biotic windows for growth of *P. putida* for eight mechanistically distinct classes of stress using 37 habitat-relevant substances (Table 2). This bacterium was able to tolerate all substances within each group of stressor; this is the first data set to demonstrate that a mesophilic bacterium can tolerate the following combination of stresses: up to 4–11 mM hexane, toluene and benzene; 8–115 mM concentrations of aromatic, chaotropic solutes; 300–825 mM chaotropic salts; 1300–2160 mM sugars or polyols; 200–750 mM kosmotropic salts; and considerable levels of matric stress (Brown, 1990) generated by kosmotropic polysaccharides

(Table 2). The *P. putida* tolerance limits to highly chaotropic substances and hydrophobic compounds were superior to those of other microbes (including extremophiles and polyextremophiles; Table 2; Hallsworth *et al.*, 2007; Bhaganna *et al.*, 2010). Whereas prokaryotes are generally less tolerant to solute stress than eukaryotes (Brown, 1990), the resilience to this array of mechanistically diverse solute stresses (Table 2) has neither been equalled nor been surpassed by any microbial species to our knowledge.

Pseudomonas putida is able to avoid contact with or prevent accumulation of stressors by using solvent-efflux pumps and synthesizing extracellular polymeric substances to form a protective barrier (Table 3). This bacterium also utilizes highly efficient macromolecule-protection systems that are upregulated in response to multiple environmental stresses including between 10 and 20 protein-stabilization proteins, oxidative-stress responses, and upregulation of energy generation and overall protein synthesis (Table 3). Furthermore, *P. putida* can synthesize an array of compatible solutes including betaine, proline, *N*-acetylglutaminylglutamine amide, glycerol, mannitol and trehalose (Table 3). Individual compatible solutes are unique in their physicochemical properties, interactions with macromolecular systems and functional efficacy for specific roles within the cell (e.g. see Chirife *et al.*, 1984; Crowe *et al.*, 1984; Brown, 1990; Hallsworth and Magan, 1994a; 1995; Yancey, 2005; Bhaganna *et al.*, 2010; Bell *et al.*, 2013). Microbes that are able to selectively utilize/deploy any substance(s) from a range of diverse compatible solutes have enhanced tolerance to diverse stresses (e.g. Hallsworth and Magan, 1995; Bhaganna *et al.*, 2010). The range of compatible solutes utilized by *P. putida* may be unique for a bacterium (see Brown, 1990) and can confer tolerance to diverse stresses including those imposed by chaotropes and hydrocarbons (Bhaganna *et al.*, 2010).

The exceptionally wide growth-windows of *Aspergillus* species in relation to a number of stress parameters correlate with their ability to dominate communities in both extreme and non-extreme (but physicochemically dynamic) environments including soils, plant necromass, saline and very high-sugar habitats (Table S1). At least 20 species of *Aspergillus* and the closely related genus *Eurotium* are xerotolerant (Pitt, 1975; Brown, 1990). *Aspergillus penicillioides* is capable of hyphal growth and conidial germination down to water activities of 0.647 and 0.68 respectively (Pitt and Hocking, 1977; Williams and Hallsworth, 2009); *Aspergillus echinulatus* can germinate down to 0.62 (Snow, 1949). Halotolerant species such as *Aspergillus wentii* can grow on the highly chaotropic salt guanidine-HCl at a higher concentration than any other microbe; up to 930 mM ($\approx 26.5 \text{ kJ g}^{-1}$ chaotropic activity; Hallsworth *et al.*, 2007) and, along with *Aspergillus*

Table 2. Growth windows for *Pseudomonas putida* under habitat-relevant stresses.^a

Type of stressor	Specific examples	Stressor concentration (mM) causing <i>P. putida</i> growth-rate inhibition of:		Notes and references
		50% ^b	100% ^b	
Hydrophobic compounds ^c	1,2,3-Trichlorobenzene ^d	0.20	0.40	<i>Pseudomonas putida</i> can be exposed to hydrophobic stressors in plant exudates, hydrocarbon-contaminated environments, and as hydrophobic catabolites and antimicrobials from <i>P. putida</i> or other microbes (see Table 6; Timmis, 2009; Bhaganna <i>et al.</i> , 2010)
	γ -Hexachlorocyclohexane ^{d,e}	0.00034	0.00072	
	2,5-Dichlorophenol ^d	0.70	0.90	
	n-hexane ^d	0.45	0.71	
	Toluene ^d	4.0	4.6	
Surfactants and aromatic solutes (highly chaotropic)	Benzene ^d	8.4	11.0	Cells come into contact with biosurfactants of microbial origin, and with chaotropic aromatics from the sources listed above (see also Hallsworth <i>et al.</i> , 2003a; Timmis, 2009; Bhaganna <i>et al.</i> , 2010)
	Tween® 80 ^{f, g}	108	164	
	Phenol ^h	9.0	13.0	
	o-cresol ⁱ	3.82	8.10	
	Benzyl alcohol ^h	20.0	32.7	
Chaotropic salts	Sodium benzoate ^f	69.2	115	Chaotropic salts occur in soils, sediments, evaporite deposits, the Dead Sea, brine lakes (including the Discovery Basin located beneath the Mediterranean Sea and the Don Juan Pond in the Antarctic) and marine-associated habitats including bitterns (see Table S1); can be highly stressful for, and lethal to, microbial systems (Hallsworth <i>et al.</i> , 2003a; 2007; Duda <i>et al.</i> , 2004)
	CaCl ₂ ^f	210	302	
	MgCl ₂ ^d	182	311	
	Guanidine-HCl ^h	125	640	
	LiCl ^h	595	825 ⁱ	
Alcohols and amides (moderately chaotropic)	Butanol ^d	158	208	Alcohols are produced by some species as antimicrobials (Table 6); ethanol is used as a biocide and can occur as an anthropogenic pollutant. Urea may be produced as an antimicrobial by some bacteria (Table 6) and is used as an agricultural fertilizer to which soil bacteria are exposed
	Ethanol ^h	600	993 ⁱ	
	Urea ^h	650	950	
	Formamide ^d	168	1080 ⁱ	
	Methanol ^d	970	1440	
Relatively neutral substances	Ethylene glycol ^h	1300	2160 ⁱ	Some organic compounds have little chao- or kosmotropic activity at biologically relevant concentrations. At low concentrations glycerol is relatively neutral but, despite its role as a stress protectant, can act as a chaotropic stressor at molar concentrations (Williams and Hallsworth, 2009)
	Glucose ^f	840	1400	
	Glycerol ^f	770	2120 ⁱ	
	Maltose ^f	326	1300	
	Proline ^f	800	2000	
Kosmotropic compatible solutes	Sorbitol ^f	503	1200	The majority of microbial compatible solutes are kosmotropic, including other compatible solutes found in <i>P. putida</i> and other bacteria such as mannitol, betaine and ectoine. <i>Pseudomonas</i> may also come into contact with some of these compounds via exposure to plant exudates
	Dimethyl sulfoxide ^f	353	1300	
	Trehalose ^f	474	771	
	Glycine ^f	99	370	
	Betaine ^f	1570	2560	
Kosmotropic salts	NaCl ^f	650	750	NaCl is environmentally ubiquitous, and is the dominant salt in most marine habitats including bioaerosols (Table S1); ammonium sulfate and KH ₂ PO ₄ are commonly applied to soils as fertilizers. In many habitats kosmotropic salts are the primary osmotic stressors, but stresses imposed on the cell are determined by the net combination of solute activities of ions and other substances present (see Hallsworth <i>et al.</i> , 2007)
	KH ₂ PO ₄ ^f	118	200	
	Ammonium sulfate ^f	198	350	
	Sodium citrate ^f	53	500	
	Dextran 40000 ^f	10	17	
Polysaccharides (kosmotropic)	Polyethylene glycol 3350 ^f	43	120	<i>Pseudomonas putida</i> is exposed to the kosmotropic activities of and/or matrix stress induced by polysaccharides such as microbe- and plant-derived extracellular polymeric substances, humic substances, etc.
	Polyethylene glycol 6000 ^f	20	40	

a. Pertinent to the phyllosphere, rhizosphere, hydrocarbon-polluted environments, and other habitats (see Table S1), including catabolic products of hydrocarbon degradation, antimicrobial substances (see Table 6), compatible solutes, and other *P. putida* metabolites. At sufficient concentrations solutes such as glucose, maltose, proline, sorbitol, glycine, betaine, NaCl and KH₂PO₄ induce osmotic stress; whereas dextran and high *M_r* polyethylene glycols induce matrix stress (see Brown, 1990); hydrophobic substances typically induce a (chaotically-mediated) hydrocarbon-induced water stress (Bhaganna *et al.*, 2010; Cray *et al.*, 2013); and other aromatics, alcohols, amides and specific salts cause chaotrope-induced water stress (Hallsworth *et al.*, 2003a; Cray *et al.*, 2013).

b. Relative to no-added stressor controls (see Hallsworth *et al.*, 2003a).

c. Log *P* > 1.95 (see Bhaganna *et al.*, 2010; Cray *et al.*, 2013).

d. These data were taken from Bhaganna and colleagues (2010); *P. putida* KT2440 (DSMZ 6125) was grown in a minimal mineral-salt broth (with glucose and NH₄Cl as the sole carbon and nitrogen substrates respectively; see Hartmans *et al.*, 1989; Bhaganna *et al.*, 2010) at 30°C. The media for control treatments had no stressors added. Some *P. putida* strains can tolerate benzene at up to 20 mM (Volkers *et al.*, 2010).

e. The pesticide γ -hexachlorocyclohexane (γ -HCH) is also known as lindane.

f. *Pseudomonas putida* KT2440 was grown in modified Luria–Bertani broth at 30°C; stressors were incorporated into media prior to inoculation, and growth rates calculated, as described by Hallsworth and colleagues (2003a). The media for control treatments had no stressors added.

g. Tween® 80 was used as a model substitute for biosurfactants (Cray *et al.*, 2013).

h. These data were taken from Hallsworth and colleagues (2003a); *P. putida* KT2440 was grown in modified Luria–Bertani broth at 30°C as described in footnote (f).

i. Extrapolated values.

j. Glycerol is a notable exception (Williams and Hallsworth, 2009).

Table 3. Stress tolerance and energy generation: cellular characteristics that contribute to weed-like behaviour.^a

Metabolites, proteins, molecular characteristics, and other traits	Function(s)	Phenotypic traits	Notes and references
Tolerances to multiple environmental stresses			
Ability to select from an array of functionally diverse compatible solutes in response to specific stress parameters	Specific compatible solutes can protect against osmotic stress, temperature extremes, freezing, dehydration and rehydration, chaotropicity, and/or hydrophobic stressors (Crowe <i>et al.</i> , 1984; Brown, 1990; Hallsworth <i>et al.</i> , 2003b; Bhaganna <i>et al.</i> , 2010; Chin <i>et al.</i> , 2010; Bell <i>et al.</i> , 2013)	Ability to maintain functionality of macromolecular systems and/or cell turgor under diverse stresses	Some microbes utilize a wide array of compatible solutes depending on the type and severity of stress; e.g. <i>Pseudomonas putida</i> is exceptional among bacteria in its ability to synthesize and accumulate protectants as diverse as betaine, proline, N-acetylglutaminylglutamine amide, glycerol, mannitol and trehalose (D'Souza-Ault <i>et al.</i> , 1993; Kets <i>et al.</i> , 1996; Bhaganna <i>et al.</i> , 2010); <i>Saccharomyces cerevisiae</i> can accumulate proline, glycerol and trehalose (see Table 6; Kaino and Takagi, 2006). Like other fungi, <i>Aspergillus</i> species utilize a range of polyols as well as trehalose (Hallsworth and Magan, 1994c; Hallsworth <i>et al.</i> , 2003b) but their exceptional ability to generate energy under stress (see below) may aid glycerol retention and thereby extending the growth window at low water activity (Hocking, 1993; Williams and Hallsworth, 2009)
Upregulation of protein-stabilization proteins in combinations tailored to specific stresses ^b	In addition to stressors (including those of biotic origin; Table 6), protein-stabilization proteins also protect against perturbations induced by stresses such as temperature extremes and desiccation (Arsene <i>et al.</i> , 2000; Ferrer <i>et al.</i> , 2003)	Maintenance of protein structure and function during environmental challenges	Several studies show highly efficient responses of <i>P. putida</i> to chaotropic and hydrophobic stressors via upregulation of between 10 and 20 protein-stabilizing proteins (Hallsworth <i>et al.</i> , 2003a; Santos <i>et al.</i> , 2004; Segura <i>et al.</i> , 2005; Domínguez-Cuevas <i>et al.</i> , 2006; Tsirogianni <i>et al.</i> , 2006; Volkers <i>et al.</i> , 2006; Ballerstedt <i>et al.</i> , 2007). The PaIA protein in <i>Aspergillus nidulans</i> induces a rapid upregulation of protein-stabilization proteins in response to extreme pH (Freitas <i>et al.</i> , 2011); high production rates of protein-stabilization proteins enhance tolerance to chaotropic salts and alcohols in <i>S. cerevisiae</i> and <i>Lactobacillus plantarum</i> , to hydrophobic stressors in <i>S. cerevisiae</i> and <i>Escherichia coli</i> (for references, see Bhaganna <i>et al.</i> , 2010), and low temperature, salt and other stresses in <i>Rhodotorula mucilaginosa</i> (Lahav <i>et al.</i> , 2004)
Intracellular accumulation of ions for osmotic adjustment; intracellular proteins structurally adapted to high ionic strength Diverse proteins and pathways conferring tolerance to solvents, chaotropic solutes and hydrophobic stressors	The utilization of ions for osmotic adjustment avoids the energy-expensive synthesis of organic compatible solutes (Kushner, 1978; Oren, 1999) Solvent-tolerant species such as <i>P. putida</i> can function at high concentrations of chaotropic and hydrophobic stressors due to the collective activities of solvent pumps that expel solvent molecules, pathways that catabolize the solvent, membrane-stabilization proteins (see below), compatible solutes (see above), and additional chaotrope- and solvent-stress responses (Ramos <i>et al.</i> , 2002; Timmis, 2002; Hallsworth <i>et al.</i> , 2003a; Volkers <i>et al.</i> , 2006; Bhaganna <i>et al.</i> , 2010; Fillet <i>et al.</i> , 2012)	Ability to maintain metabolic activity at high extra- and intracellular ionic strength High tolerance to chaotropic and hydrophobic solvent stressors	<i>Halobacterium salinarum</i> (unusually for a bacterium), utilize the so-called 'salt-in' strategy (Oren <i>et al.</i> , 2002; Bardavid and Oren, 2012; Saum <i>et al.</i> , 2012) Due to its exceptional solvent tolerance <i>P. putida</i> is the organism of choice for many industrial applications including biotransformation in two-phase systems and bioremediation (Timmis, 2002).
Membrane-stabilization proteins: e.g. <i>cis/trans</i> -isomerase (CTI)	Membrane stabilization under stress	Enhanced tolerance to stresses that impact membrane structure	Several proteins are involved in maintaining bilayer rigidity in <i>P. putida</i> during exposure to toluene and other solvents or hydrophobic stressors (Bernal <i>et al.</i> , 2007)
Proteins involved in the moderation of membrane fluidity	Maintenance of membrane fluidity and membrane-associated processes under conditions such as low temperature (Turk <i>et al.</i> , 2011)	Enhanced tolerance to diverse, physicochemical stress parameters	For example <i>Aureobasidium pullulans</i> , <i>Cryptococcus liquefaciens</i> and <i>R. mucilaginosa</i> (Turk <i>et al.</i> , 2011)

Promoter(s) that simultaneously control multiple stress-response genes, e.g. stress-response elements (STREs)	Activation of multiple cellular stress-responses	Enhanced tolerance to multiple stress-parameters	The transcription factors (Msn2p and Msn4p) involved in STRE-mediated stress responses in <i>S. cerevisiae</i> accumulate in the nucleus upon heat-, ethanol-, (chaotrope-), and osmotically induced stresses, and during nutrient limitation. The STRE is upstream of genes for protein-stabilization proteins (see above) metabolic enzymes, and cellular organization proteins (Görner <i>et al.</i> , 1998; Moskvina <i>et al.</i> , 1998)
Proteins involved in DNA protection under stress; e.g. DNA-binding protein (Dps)	Formation of a protein:chromosome complex that protects DNA	Enhanced DNA stability and reduced risk of mutation in relation to multi-farious stresses	The role of Dps has been well-characterized in <i>E. coli</i> in relation to diverse environmental challenges including reactive oxygen species, iron and copper toxicity, thermal stress and extremes of pH (Nair and Finkel, 2004)
Pathways for synthesis of carotenoids (e.g. β -carotene in <i>Dunaliella salina</i> ; torularhodin in <i>Rhodotorta</i> spp.)	Carotenoids absorb light and thereby protect cellular macromolecules and organelles from exposure to ultraviolet	Tolerance of exposure to ultraviolet irradiation	Synthesis of β -carotene is upregulated in response to light in <i>D. salina</i> (Chen and Jiang, 2009); carotenoids are concentrated in the plasma membrane of <i>S. ruber</i> (Antón <i>et al.</i> , 2002); studies of <i>R. mucilaginosa</i> show that torularhodin enhances tolerance to ultraviolet B exposure (Moliné <i>et al.</i> , 2010)
Preferential utilization of a chaotropic compatible solute at high NaCl concentration and/or low temperature	Rigidification of macromolecular structures at high NaCl and low temperature can induce metabolic inhibition (Brown, 1990; Ferrer <i>et al.</i> , 2003; Chin <i>et al.</i> , 2010) that can be compensated for by glycerol and/or fructose (Brown, 1990; Chin <i>et al.</i> , 2010)	Enhanced tolerance to extreme conditions that rigidity macromolecular systems	The high intracellular concentrations of glycerol in <i>D. salina</i> under salt stress and glycerol and/or fructose in and <i>Aspergillus</i> and <i>Eurotium</i> spp. and <i>R. mucilaginosa</i> at low temperature are able to extend growth windows under these extreme conditions (Brown, 1990; Lahav <i>et al.</i> , 2002; Chin <i>et al.</i> , 2010). <i>Dunaliella salina</i> has membranes with reduced glycerol permeability thereby aiding retention of this compatible solute (Bardavid <i>et al.</i> , 2008)
Efficient system to eliminate free radicals (see above)	Elimination of reactive oxygen species (see also Table 5)	Various substances and stress parameters can induce oxidative stress	Diverse environmental substances and parameters induce oxidative stress, either directly or indirectly (e.g. via lipid peroxidation). For example, chaotropic stressors and heat shock have been associated with the induction of oxidative stress responses in <i>S. cerevisiae</i> and <i>P. putida</i> (Costa <i>et al.</i> , 1997; Hallsworth <i>et al.</i> , 2003a)
Proteins associated with regulation of cytosolic pH; e.g. the lysine decarboxylase system (coded for by the <i>cadBA</i> operon); the Na ⁺ /H ⁺ antiporter	Conversion and export of lysine/cadaverine that results in increased pH of the cytosol in bacteria (Álvarez-Ordóñez <i>et al.</i> , 2011); the Na ⁺ /H ⁺ antiporter Nha1 decreases intracellular hydrogen ion concentration	Tolerance of low-pH environments	Decarboxylase systems that increase intracellular pH also exist for other amino acids (Álvarez-Ordóñez <i>et al.</i> , 2011); Nha1 greatly enhances viability of <i>Saccharomyces boulardii</i> at pH 2 (dos Santos Sant'Ana <i>et al.</i> , 2009)
Multidrug efflux pumps (for details see Table 6) Enzymes for synthesis of extracellular polymeric substances	Production and secretion of polymeric substances	Enhanced tolerance to multiple stressors and stress parameters	Extracellular polymeric substances have been shown to confer protection to <i>E. coli</i> cells exposed to high temperature, low pH, salt and oxidative stress (Chen <i>et al.</i> , 2004), and are associated with stress protection in <i>Clostridium</i> spp., <i>Pantoea ananatis</i> , <i>P. putida</i> and other <i>Pseudomonas</i> spp. (Chang <i>et al.</i> , 2007; Morohoshi <i>et al.</i> , 2011; see also Table 6)
Upregulation of proteins involved in protein synthesis under stress	To enhance or maintain the quantity, type, and functionality of cellular proteins under stress	Maintenance of the cellular system and its metabolic activity under stress Enhanced stress tolerance	Can be upregulated in response to stress induced by high temperature, chaotropic solutes, hydrophobic stressors and other stress parameters (e.g. <i>P. putida</i> ; Hallsworth <i>et al.</i> , 2003a; Volkers <i>et al.</i> , 2009)
Polyplidity, aneuploidy, association with plasmids, and/or hybridization that enhance(s) stress tolerance	Whereas hybridization, polyplidity, aneuploidy and association with plasmids can be ecologically disadvantageous, such traits can enhance the vigour, stress tolerance and competitive ability of some microbes such as <i>Pichia</i> spp., <i>S. cerevisiae</i> , <i>P. putida</i> and <i>S. ruber</i> (Table 1; Rainieri <i>et al.</i> , 1999; Naumov <i>et al.</i> , 2001; Mongodin <i>et al.</i> , 2005; Tark <i>et al.</i> , 2005; Dhar <i>et al.</i> , 2011; Chen <i>et al.</i> , 2012)		Aneuploid cells of <i>S. cerevisiae</i> have enhanced tolerance to oxidative stress and diverse inhibitors of biochemical processes (Chen <i>et al.</i> , 2012); polyplidity can enhance vigour and stress tolerance in some species (see also Tables 1 and 5); the stress tolerance of some microbial weeds is enhanced by their association with plasmids (e.g. the <i>P. putida</i> tol-plasmid that enhances solvent tolerance; Tark <i>et al.</i> , 2005; Domínguez-Cuevas <i>et al.</i> , 2006; the <i>S. ruber</i> plasmid contains a gene involved in ultraviolet protection; Mongodin <i>et al.</i> , 2005)

Table 3. *cont.*

Metabolites, proteins, molecular characteristics, and other traits	Function(s)	Phenotypic traits	Notes and references
Polyextremophilic growth-phenotype	Ability to grow optimally under multiple, extreme conditions; e.g. extremely acid and alkali (pH 2–10), high-salt (2.5 M NaCl), and low-temperature conditions (Lahav <i>et al.</i> , 2002; Libkind and Sampalo, 2010)	Polyextremophile	<i>Rhodotomila mucilaginosa</i> is capable of growth over an incredible range of physicochemical conditions (including almost the entire pH range for microbial life) and is able to colonize a correspondingly wide diversity of environments (for examples see main text)
Genes associated with salt stress	Sugar catabolism (e.g. <i>RmPGM2</i> , <i>RmGPA2</i> , <i>RmACK1</i> , <i>RmMCP1</i> and <i>RmPET9</i>), protein glycosylation (e.g. <i>RmSEC53</i>), aromatic amino acid synthesis (e.g. <i>RmARO4</i>), regulation of protein synthesis (e.g. <i>RmANB1</i>)	Maintenance of protein synthesis and energy generation under salt stress	Between 10 and 20 genes are involved in increased tolerance to NaCl and/or LiCl in <i>R. mucilaginosa</i> (Lahav <i>et al.</i> , 2004; Gostinčar <i>et al.</i> , 2012)
Metabolites associated with resistance to antimicrobial plant metabolites and survival in phyllosphere habitats	Metabolites involved in quorum sensing may enhance competitive ability on the leaf surface	Enhanced survival and competitive ability	By contrast to other phyllosphere microbes, <i>P. ananatis</i> is resistant to the inhibitory effects of plant alkaloids and this correlates with the production of metabolites involved in quorum sensing such as <i>N</i> -acyl-homoserine lactone (Enya <i>et al.</i> , 2007)
High-efficiency energy-generation systems Light-activated proton pumps (e.g. xanthorhodopsin)	To generate a proton-motive force, utilizing light as an energy source	Utilization of light for transmembrane proton transport	Xanthorhodopsin, found in <i>S. ruber</i> , is a proton pump made up of two chromophores; a light-absorbing carotenoid and a protein (Balashov <i>et al.</i> , 2005); a light-activated proton pump was also recently described in <i>H. walsbyi</i> (Lobasso <i>et al.</i> , 2012)
High expression of genes involved in energy generation (e.g. <i>fta</i> , <i>pykF</i> , <i>atpA</i> and <i>atpD</i>)	Rapid catabolism of carbon substrates leading to enhanced synthesis of ATP	Efficient energy-generation	High expression of <i>E. coli</i> genes <i>fta</i> , <i>pykF</i> , <i>atpA</i> and <i>atpD</i> can be associated with ability to grow rapidly, enhanced stress tolerance and – by implication – greater competitive ability (Karlin <i>et al.</i> , 2001)
Enhanced energy-efficiency at low nutrient concentrations (e.g. see glutamate dehydrogenase entry in Table 5)	Maintenance of energy generation close to the point of system failure under hostile conditions	Efficient energy generation during potentially lethal challenges	The exceptional ability of <i>P. putida</i> to respond and adapt to chaotropic, solvent and hydrocarbon stressors has, in part, been attributed to considerable increases in energy- and NAD(P)H-generating systems
Upregulation of proteins involved in energy generation under chaotropic and solvent-induced (hydrocarbon) stresses (Hallsworth <i>et al.</i> , 2003a; Volkens <i>et al.</i> , 2006)	Enhanced primary metabolism and energy-generation systems	Enhanced vigour and energy production	<i>Aspergillus</i> species have numerous genome modifications, such as gene duplications for pyruvate dehydrogenase, citrate synthase, aconitase and malate dehydrogenase that enhance primary metabolism including glycolysis and the TCA cycle (Filiphi <i>et al.</i> , 2009)
Isoenzymatic genome modifications associated with reinforced energy metabolism	Catabolism of stored substances to boost growth, stress tolerance, production of secondary metabolites, motility and/or other key cellular functions under stress	Ability to maintain energy production during nutrient limitation and under hostile conditions	Efficient storage (see Table 5) and utilization of energy-generating substances such as trehalose, glycogen and polyhydroxyalkanoates confers a competitive advantage in <i>S. cerevisiae</i> and other species (Castrol <i>et al.</i> , 1999; Kadouri <i>et al.</i> , 2005)
Efficient utilization of intracellular reserves for energy generation	Via aerobic fermentation <i>S. cerevisiae</i> can simultaneously generate energy while synthesizing the chaotropic stressor ethanol (see Table 6)	Energy generation linked to synthesis of an antimicrobial	Each glucose molecule fermented by <i>S. cerevisiae</i> produces two ethanol molecules and two molecules of ATP (Piskur <i>et al.</i> , 2006)
Concomitant energy generation and synthesis of antimicrobials with no loss of ATP-generation efficiency			

a. The current table provides examples of proteins, genes or other characteristics that can be associated with and give rise to weediness; the categories of traits are not mutually exclusive. Individual traits may not be unique to microbial weeds; however, weeds species are likely to have a number of these types of characteristics.

b. For example chaperonins, heat-shock proteins and cold-shock proteins.

c. That code for fructose-1,6-bisphosphate aldolase, pyruvate kinase and ATP synthase subunits F1 α and F1 β respectively.

oryzae, up to ~ 4 M NaCl or KCl (Hallsworth *et al.*, 1998). *Aspergillus* (and *Eurotium*) strains are capable of reasonable growth rates close to 0°C (Chin *et al.*, 2010), and some reports suggests that the biotic windows for some *Aspergillus* species ultimately fail at $\leq -18^{\circ}\text{C}$ (Vallentyne, 1963; Siegel and Speitel, 1976). Whereas a number of comparable fungal genera show moderate levels of stress tolerance, *Aspergillus* species generally out-perform their competitors and top numerous league tables for ascomycete stress tolerance (see Pitt, 1975; Wheeler and Hocking, 1993; Williams and Hallsworth, 2009).

Studies of seven *Aspergillus* species have revealed a number of genome modifications within this genus that reinforce both primary metabolism and energy generation (Table 3; Flipphi *et al.*, 2009). These include gene duplications for key enzymes that control metabolic flux, such as pyruvate dehydrogenase, citrate synthase, aconitase, and malate dehydrogenase that enhance glycolysis and the TCA cycle (Flipphi *et al.*, 2009). There is evidence that, at the low water activities (high osmotic pressures) associated with very high-salt and high-sugar habitats, the biotic window of xerophilic fungi fail due to the prohibitive energy expenditure required to prevent the high levels of intracellular glycerol from leaking out of the membrane (Hocking, 1993). The adaptability and competitive ability of *Aspergillus* species (most of which can also proliferate under non-extreme conditions) may be attributed, in part, from enhanced energy-generating capability. *Aspergillus* species (like many fungi) can utilize a range of polyols – as well as trehalose – as compatible solutes based on their individual strengths as protectants against osmotic stress, desiccation rehydration, chaotropes etc. (Crowe *et al.*, 1984; Brown, 1990; Hallsworth *et al.*, 2003b). *Aspergillus* and *Eurotium* strains can also accumulate chaotropic sugars from the medium, or preferentially synthesize and accumulate chaotropic glycerol, in order to enhance growth at temperatures below ~ 10°C (Chin *et al.*, 2010; see also below).

Salinibacter ruber is a halophilic bacterium found in high numbers at molar salt concentrations in aerobic aquatic habitats that are otherwise more densely populated by archaeal than bacterial species. Unusually for a bacterium, *S. ruber* takes in ions to use as compatible solutes and thereby avoids synthesis of organic compatible solutes that would be energetically unfavourable in environments of > 1 M NaCl (Oren, 1999) and has evolved intracellular proteins that are both structurally stable and catalytically functional in the cytosol at high ionic strength (Table 3). *Salinibacter ruber* (as well as the alga *Dunaliella salina* and yeast *R. mucilaginosa*) accumulates carotenoids in response to light which can protect against oxidative stress that is a potential hazard in salt-erns exposed to high levels of ultraviolet (Table 3). *Haloquadratum walsbyi*, an Archaeon that is also a dominant

player in saltern communities (Table S1), also uses the 'salt-in' compatible-solute strategy, and – along with *S. ruber* – has light-activated protein pumps that generate proton-motive force and thereby boost energy generation (Table 3). In addition, *H. walsbyi* is unusually tolerant to the chaotropic salt MgCl_2 that can reach high concentrations in evaporate ponds (Hallsworth *et al.*, 2007; Bardavid *et al.*, 2008). *Dunaliella salina* is also highly prevalent in saline habitats that (unusually for an alga) can successfully compete with the multitude of halophilic Archaea found in 5 M NaCl environments (see later; Table S1). This alga preferentially utilizes glycerol as a compatible solute, is able to accumulate glycerol to molar concentrations (6–7 M; see Bardavid *et al.*, 2008), and has low membrane permeability to this compatible-solute thus aiding retention (Bardavid *et al.*, 2008).

Glycerol is unique in its ability to maintain the flexibility of macromolecular structures under conditions that would otherwise cause excessive rigidification, e.g. molar concentrations of NaCl, or low temperature (Back *et al.*, 1979; Hallsworth *et al.*, 2007; Williams and Hallsworth, 2009; Chin *et al.*, 2010). Species able to grow at low or subzero temperatures (e.g. *Aspergillus*, *Cryptococcus*, *Rhodotorula* species, including *R. mucilaginosa*) either primarily use glycerol as a compatible solute or preferentially accumulate this chaotrope at low temperature (Table 3; Lahav *et al.*, 2004; Chin *et al.*, 2010). *Rhodotorula mucilaginosa* is able to colonize – and frequently dominates – diverse substrates/environments including saline and non-saline soils, aquatic habitats (both liquid- and ice-associated), the acidic surface-film of sphagnum moss (as well as other plant tissues) and various surfaces within the human body and other animals (Table S1; Egan *et al.*, 2002; Trindade *et al.*, 2002; Vital *et al.*, 2002; Lahav *et al.*, 2002; 2004; Lisichkina *et al.*, 2003; Buzzini *et al.*, 2005; Perniola *et al.*, 2006; Goyal *et al.*, 2008; Turk *et al.*, 2011; Kachalkin and Yurkov, 2012). This ability is coupled with an exceptional level of tolerance to multiple stresses: *R. mucilaginosa* is (highly unusual for a yeast) a polyextremophile that is psychrophilic, highly salt-tolerant, and can growth over the majority of the pH range known to support microbial life (i.e. < 2 and > 10, see Table 3; Lahav *et al.*, 2002).

Some microbial habitats such as fruits that have been disconnected from the vascular system (and immune responses) of the parent plant have a favourable water activity and high nutrient availability and are therefore potentially habitable by a wide range of microbes (Ragaert *et al.*, 2006). As the microbial community develops on a detached fruit (or in apple or grape musts; see Table S1) there can be changes in the composition and concentration of sugars, osmolarity, water activity, ethanol concentration (and that of other antimicrobials), pH and other environmental factors that are severe enough to

Fig. 2. Growth of *Saccharomyces cerevisiae* strains: (A) CCY-21-4-13, (B) Alcotec 8 and (C) CBS 6412 and other yeast species: (D) *Candida etchellsii* (UWOPS 01-168.3), (E) *Cryptococcus terreus* (PB4), (F) *Debaryomyces hansenii* (UWOPS 05-230.3), (G) *Hansenula (Ogataea) polymorpha* (CBS 4732), (H) *Hortaea werneckii* (MZKI B736), (I) *Kluyveromyces marxianus* (CBS 712), (J) *Pichia (Kodamaea) ohmeri* (UWOPS 05-228.2), (K) *Pichia (Komagataella) pastoris* (CBS 704), (L) *Pichia sydowiorum* (UWOPS 03-414.2), (M) *Rhodotorula creatinivora* (PB7), (N) *Saccharomycodes ludwigii* (UWOPS 92-218.4) and (O) *Zygosaccharomyces rouxii* (CBS 732) on a range of media at 30°C. These were: malt-extract, yeast-extract phosphate agar (MYPIA) without added solutes (control), and MYPIA supplemented with diverse stressors – ethanol, methanol, glycerol, fructose, sucrose, NaCl, MgCl₂, ammonium sulfate, proline, xanthan gum and polyethylene glycol (PEG) 8000 – over a range of concentrations as shown in key (values indicate %, w/v). A standard Spot Test was carried out (see Chin *et al.*, 2010; Toh *et al.*, 2001) that was modified from Albertyn and colleagues (1994) for stress phenotype characterization (see Table 5). Colony density was assessed after an incubation time of 24 h on a scale of 0–5 arbitrary units (Chin *et al.*, 2010). Cultures were obtained from (for strain A) the Culture Collection of Yeasts (CCY, Slovakia); (strain B) Hambleton Bard Ltd, Chesterfield, UK; (strains C, G, I, K, O) the Centraalbureau voor Schimmelcultures (CBS, the Netherlands); (strains D, F, J, L, N) the University of Western Ontario Plant Sciences Culture Collection (UWOPS, Canada); (strains E, M) were obtained from Dr Rosa Margesin, Institute of Microbiology, Leopold Franzens University, Austria; and (strain H) the Microbial Culture Collection of National Institute of Chemistry (MZKI, Slovenia). All Petri plates containing the same medium were sealed in a polythene bag to maintain water; all experiments were carried out in duplicate, and plotted values are the means of independent treatments.

inhibit or eliminate most members of the original microflora (see Brown, 1990; Hallsworth, 1998; de Pina and Hogg, 1999; Pretorius, 2000; Bhaganna *et al.*, 2010). The yeast *S. cerevisiae* that is relatively scarce in natural habitats (and even when present forms a quantitatively insignificant component of the natural community; Renouf *et al.*, 2005; Raspor *et al.*, 2006) is unsurpassed in its ability to emerge to dominate high-sugar habitats at water activities above 0.9 (Table S1; Hallsworth, 1998; Pretorius, 2000). We therefore compared its tolerance towards a range of habitat-relevant stressors with those of 12 other yeasts that occur at the insect–flower interface, on plant tissues, and/or are environmentally widespread (Fig. 2; Table 4). This experiment revealed that *S. cerevisiae* has a more vigorous growth phenotype and more robust stress tolerance than the other species tested with the exception of two species of *Pichia*, another genus characterized by weed-like traits (see Fig. 2; Table 4). Stress phenotypes of the yeasts assayed were compared on the basis of stressor type and mechanistically distinct stress-parameters in order to assess resilience to both the degree and diversity of environmental challenges (Table 4). Under the assay conditions it was the three *S. cerevisiae* strains, *Pichia (Kodamaea) ohmeri* and *Pichia sydowiorum* that exhibited the greatest tolerance to the widest range of solute-imposed stresses (Table 4). Other species studied were slow growers, have specific environmental requirements (e.g. the halotolerant *D. hansenii* and *Hortaea werneckii*), and/or otherwise lacked the capacity for vigorous growth on the high-sugar substrate at 30°C (see Fig. 2). Generally *S. cerevisiae* strains were fast-growing, xerotolerant, and highly resistant to chaotrope-induced, osmotic, matric and kosmotrope-induced stresses and are known to be acidotolerant and capable of growth over a wide temperature range, from ~0°C to 45°C (Fig. 2A and B; Table 4). Although *Pichia* species can produce ethanol (*P. anomala* can produce up to 0.36 M ethanol; Djelal *et al.*, 2005) when growing side by side in high-sugar habitats with *S. cerevisiae*, *Pichia* species are unable to tolerate the

molar ethanol concentrations that *S. cerevisiae* can produce (Lee *et al.*, 2011).

Saccharomyces cerevisiae is metabolically wired to concomitantly synthesize an antimicrobial chaotrope (ethanol), accumulate a compatible solute (glycerol) that affords protection against this chaotrope (see below), and at the same time generate energy without any loss of ATP-generation efficiency (Table 3): a remarkable phenotype that is coupled with the deployment of other compatible solutes under specific conditions (trehalose and proline). Trehalose, which can reach concentrations of up to 13% dry weight in *S. cerevisiae* (Aranda *et al.*, 2004), is highly effective at stabilizing macromolecular structures that are exposed to chaotropic substances (e.g. ethanol, 2-phenylethanol) and enabling cells to survive desiccation–rehydration cycles (Crowe *et al.*, 1984; Mansure *et al.*, 1994). The highest reported tolerance to ethanol stress was in *S. cerevisiae* cells capable of growth and metabolism up to 28% v/v (\equiv 20% w/v or 4.3 M; chaotropic activity = 25.3 kJ g⁻¹; Hallsworth, 1998; Hallsworth *et al.*, 2007) when the medium contained high concentrations of a compatible solute, proline (Thomas *et al.*, 1993); furthermore of the seven microbes reported to have the highest resistance to other chaotropic substances (including urea, phenol) the majority are weed species including *A. wentii*, *S. cerevisiae*, *Escherichia coli* and *P. putida* (Hallsworth *et al.*, 2007).

The stress mechanisms for numerous types of cellular stress operate at the level of water:macromolecule interactions (Crowe, *et al.*, 1984; Brown, 1990; Sikkema *et al.*, 1995; Casadei *et al.*, 2002; Ferrer *et al.*, 2003; Hallsworth *et al.*, 2003a; 2007; McCammick *et al.*, 2009; Bhaganna *et al.*, 2010; Chin *et al.*, 2010). As front-line stress protectants, compatible solutes (within limits) can therefore mollify stress the inhibitory effects of an indefinite number of stress parameters (see Crowe, *et al.*, 1984; Brown, 1990; Hallsworth *et al.*, 2003b; Bhaganna *et al.*, 2010; Chin *et al.*, 2010; Bell *et al.*, 2013) but cells also upregulate other protection systems (see also above; Table 3). In *S. cerevisiae*, an extensive number of genes that code for




Table 4. Stress phenotypes and ecophysiological profiles of *Saccharomyces cerevisiae* relative to those of other yeast species.

Yeast species ^a	Tolerance to diverse stressors ^b					Tolerance to distinct stress mechanisms ^b			Additional notes on stress and growth phenotypes ^c	Typical habitat	Ecophysiological description ^d
	Alcohols	Glycerol	Sugars	Salts	Others	Chaotropic	Osmotic	Matric	Kosmotropic		
<i>Saccharomyces cerevisiae</i> strain											
CCY-21-4-13	Higher	Mid	Higher	Higher	Higher	Higher	Higher	Higher	Higher	Fruits; other fleshy plant tissues; nectar	Multiple stress tolerances including sugar and ethanol ^f ; stress phenotype of a xerotolerant, acidotolerant, vigorous weed
Abotec 8	Higher	Mid	Mid	Mid	Higher	Higher	Mid	Higher	Mid	As above	As above
CBS 6412	Higher	Mid	Higher	Mid	Higher	Higher	Mid	Higher	Mid	As above	As above
Other species (strain)											
<i>Candida etchellsii</i> (UWOPS 01-168.3)	Lower	Higher	Mid	Higher	Mixed	Higher	Higher	Mid	Mid	Pollinating insects; flowers; fermented foods	Intolerant to specific stressors; slow-growing
<i>Cryptococcus terreus</i> (PB4)	Mid	Mid	Higher	Mid	Higher	Mid	Mid	Higher	Mid	Soils and sediments; subsurface environments	A slow-growing psychrophile ^g
<i>Debaryomyces hansenii</i> (UWOPS 05-230.3)	Mid	Mid	Higher	Higher	Higher	Mid	Mid	Higher	Lower	High-salt environments and foods	Halotolerant; a fast-growing high-salt specialist
<i>Hansenula (Ogataea) polymorpha</i> (CBS 4732)	Higher	Mid	Mid	Mid	Higher	Higher	Lower	Higher	Higher	Soil, insect gut, fruit juice	Neither fast-growing nor highly stress tolerant; a high-temperature specialist
<i>Hortaea werneckii</i> (MZK1 B736)	Mid	Mid	Higher	Higher ^h	Lower	Mid	Mid	Lower	Mid	Salty environments; animal and plant surfaces	Intolerance to matric stress; a slow-growing halophile
<i>Kluyveromyces marxianus</i> (CBS 712)	Higher	Lower	Mid	Lower	Higher	Mid	Lower	Higher	Lower	Milk; plant surfaces; natural fermentations	Thermotolerant, solute-intolerant and xero-intolerant ⁱ
<i>Pichia (Kodamaea) ohmeri</i> (UWOPS 05-228.2)	Higher	Mid	Higher	Higher	Higher	Higher	Higher	Higher	Mid	Beetles; fruits and other plant	Multiple stress tolerances including ethanol and sugar; stress phenotype of a xerotolerant, vigorous weed
<i>Pichia (Komagataella) pastoris</i> (CBS 704)	Mid	Lower	Lower	Mid	Higher	Mid	Lower	Higher	Mid	Rotting wood; slimes	Intolerant to osmotic stress; glycerol and sugar; a xero-intolerant yeast
<i>Pichia sydneyorum</i> (UWOPS 03-414.2)	Higher	Mid	Higher	Higher	Higher	Higher	Mid	Higher	Higher	Polluted environments; waste water	Multiple stress tolerances including ethanol and sugar; stress phenotype of a xerotolerant, vigorous weed
<i>Rhodotorula creatinivora</i> (PB7)	Lower	Lower	Mid	Mid	Lower	Lower	Lower	Mid	Higher	Subsurface; soil	Intolerant to alcohols; glycerol; xerotolerant
<i>Saccharomycodes ludwigii</i> (UWOPS 92-218.4)	Lower	Mid	Higher	Lower	Lower	Lower	Lower	Lower	Lower	Grapes; plant tissues and fermentations	A psychrophilic slow grower
<i>Zygosaccharomyces rouxii</i> (CBS 732)	Mid	Higher	Higher	Lower	Mixed	Higher	Mid	Mid	Mid	Fruits and other high-sugar habitats	Intolerant to glycerol and NaCl; tolerant to sugars; an osmophile

^a Sources of strains are given in Fig. 2.

^b Based on data for growth on MYPIA at 30°C (see Fig. 2). Alcohols were ethanol and methanol, sugars were fructose and sucrose, salts were NaCl, MgCl₂ and ammonium sulfate, the other stressors were proline, xanthan gum and PEG 8000 (Fig. 2). Chaotropic stress is induced by ethanol, fructose, glycerol and MgCl₂; osmotic stress by fructose, sucrose, NaCl and proline; matric stress by xanthan gum and PEG 8000; and kosmotropic stress by ammonium sulfate (Fig. 2; Brown, 1990; Hallsworth, 1998; Hallsworth *et al.*, 2003a; Hallsworth *et al.*, 2007; Williams and Hallsworth, 2009; Bhaganna *et al.*, 2010; Chin *et al.*, 2010; Cray *et al.*, 2013). Stress-tolerance assessments (higher, mid-range, lower) are relative to the range of those of the other species assayed, and based on values in Fig. 2.

^c Temperature optima, minima and/or maxima (T_{opt} , T_{min} , T_{max}) and pH optima, minima and/or maxima (pH_{opt} , pH_{min} , pH_{max}) for growth were obtained from cited publications. Extreme growth phenotypes (fast or slow growers) were listed according to growth rates on MYPIA at 30°C (see also Fig. 2). Species designated as *slow growers* did not have a vigorous growth phenotype under these conditions, i.e. all strains listed are able to grow rapidly within the assay period on/at their favoured media/temperature. The designation *slow growing* did not relate to an extended lag phase (data not shown).

^d Primarily based on data presented in the current table as well as Fig. 2 (unless otherwise stated) so is most pertinent to high-sugar habitats at -30°C. Habitat dominance will ultimately be influenced by antimicrobial activities of species present within a community; see Table 6). Ecophysiological descriptions of weed-like species are underlined in bold.

^e Values for temperature- and pH-tolerances were taken from: (for *S. cerevisiae*) Praphalung and Fleet (1997), Yalcin and Ozbas (2008) and Salvador and colleagues (2011); (for *C. terreus*) Krallish and colleagues (2006); (for *D. hansenii*) Dominguez and colleagues (1997); (for *H. polymorpha*) Levine and Cooney (1973); (for *H. werneckii*) Diaz Muñoz and Montalvo-Rodriguez (2005); (for *K. marxianus*) Aziz and colleagues (2009); Salvador and colleagues (2011); (for *P. ohmeri*) Zhu and colleagues (2010); (for *P. pastoris*) Chiruvolu *et al.* (1998); (for *P. sydneyorum*) Kanekar and colleagues (2009); (for *R. creatinivora*) Krallish and colleagues (2006); (for *S. ludwigii*) Stratford (2006); and (for *Z. rouxii*) Restaino and colleagues (1983) and Praphalung and Fleet (1997). Tolerance windows for *Candida etchellsii* are yet to be established.

^f Under some conditions strains of *S. cerevisiae* can tolerate up to 28% v/v (20% w/v) ethanol (see Hallsworth, 1998).

^g See also Krallish and colleagues (2006).

^h In high-salt habitats, *H. werneckii* can grow at molar concentrations of NaCl (see Gunde-Cimerman *et al.*, 2000) but was less salt tolerant under the conditions of this assay.

ⁱ This species appears to have a requirement for high water activity.

^j Habitat dominance can also depend on factors such as production of antimicrobials (see also Table 6). *Pichia ohmeri* can dominate some habitats, *Pichia anomala* may be more vigorous as a weed species (see Rosa *et al.*, 1990; Lee *et al.*, 2011).

protein-stabilizing proteins, metabolic enzymes and cellular-organization proteins are regulated by a single promoter – the stress response element (STRE) – which is activated under cellular stress. This regulatory system enables the coordination of an array of stress responses in an energy-efficient manner (Table 3). Numerous studies show that the *S. cerevisiae* responses to osmotic stress [via the high-osmolarity glycerol-response (HOG) pathway], oxidative stress, and challenges to protein and membrane stability are exceptionally efficient (Table 3; Brown, 1990; Costa *et al.*, 1997; Hallsworth, 1998; Rodríguez-Peña *et al.*, 2010; Szopinska and Morsomme, 2010). In addition to these stress responses, there is evidence that the dynamics of polyphosphate and glycogen metabolism in *S. cerevisiae* are wired in such a way to boost ATP production (Castrol *et al.*, 1999), and that the ploidy levels of many strains enhance tolerance towards oxidative and other stresses (Table 3). The energy-generation and energy-conservation strategies employed by *S. cerevisiae* have apparent parallels in other weed species; for example, analyses of synonymous codon bias in the genomes of fast-growing bacteria including *E. coli* and *Bacillus subtilis* suggest that genes associated with energy generation are highly expressed (Karin *et al.*, 2001; Table 5; see also above). The enhanced energy efficiency that is apparent in some weed species not only underpins their robust stress biology, but is required to sustain a vigorous growth phenotype and/or to out-grow competitors.

Growth, nutritional versatility, and resource acquisition and storage

Efficient cell division and an ability to out-grow competitors are implicit to the emergence of weed species. Species that grow relatively slowly or remain quiescent, and those with specific requirements for a host, a symbiotic partner, or obligate interactions with other microbial species are disadvantaged during community development in open habitats: dominance ultimately results from vigour and – in many habitats – a degree of nutritional versatility (Tables S1 and 5). Ability to out-grow competitors is equally important (possibly even more so) for weed species in nutrient-poor and physicochemically extreme habitats that can both select for slow-growing species and limit overall biomass development. Many traits that give rise to an efficient growth phenotype can be found in weed species (Table 5): multiple copies of ribosomal DNA operons that can enhance rates of protein synthesis; transport proteins to scavenge macromolecules thus negating the need for *de novo* synthesis; high-efficiency DNA polymerase for rapid genome replication; efficient systems to eliminate reactive oxygen species that are a by-product of rapid growth; and poly-

ploidy, aneuploidy and plasmids that can be associated with enhanced stress tolerance, higher growth-rates and increased overall vigour (Table 5). We believe that another key aspect of weed biology for some microbial species is their ability to sequester and store available nutrient and substrate molecules with a high efficiency as an auxiliary energy reserve (see below), to be able to supply daughter cells with nutrients, and simultaneously deprive other microbes of nutritional resources (e.g. Duiff *et al.*, 1993; Castrol *et al.*, 1999; Loaces *et al.*, 2011; Renshaw *et al.*, 2002; Jahid *et al.*, 2006). Microbial weeds are highly efficient at nutrient-sequestration and substrate utilization, as well as intracellular storage of these substances (Table 5). For example, fungal siderophores that remove iron from the habitat have been shown to suppress growth of other microorganisms due to iron limitation (Renshaw *et al.*, 2002).

Intermittent nutrient availability and low nutrient concentrations are characteristic of many aquatic environments so the ability to scavenge and store, circumvent the requirement for, or strategically utilize scarce and potentially limiting nutrients can greatly enhance the competitive ability of planktonic species. Under phosphate-limiting conditions some aquatic species, including *S. ruber*, are able to use sulfur-containing glycolipids in place of phospholipids during low phosphate availability (Table 5). The cyanobacterium *M. aeruginosa* has high-efficiency phosphate-transport systems and accumulates intracellular polyphosphate which is used to maintain activity once extracellular phosphate is depleted (Table 5; Shi *et al.*, 2003). It is likely that this trait contributes to its weed-like phenotype in Lake Taihu (China) that is a well-studied, phosphate-limited habitat frequently dominated by *M. aeruginosa* (Shi *et al.*, 2003). Iron is another potentially limiting nutrient in aquatic habitats. *Dunaliella salina* utilizes a highly efficient iron-binding protein to take up iron which is then stored in vacuoles (Paz *et al.*, 2007). Species such as *D. salina* and *H. walsbyi* cope with oligotrophic conditions by hyperaccumulating starch and/or lipids such as polyhydroxybutyrate especially when nutrients other than carbon are limiting (Table 5; Bardavid *et al.*, 2008). One weed species with adaptations to endure low-nitrogen conditions is *Mycobacterium smegmatis*. This species (as well as other mycobacteria) which can be prevalent in relatively low-nutrient aquatic and non-aquatic habitats (Koch, 2001; Kazda and Falkinham III, 2009) have biochemical adaptations to enable growth and metabolic activity even when nitrogen concentrations are an order of magnitude lower than those usually required for structural growth (i.e. a C:N ratio of 8:1; Roels, 1983; Behrends *et al.*, 2012). In addition to nitrogen oligotrophy, *M. smegmatis* has a DNA polymerase capable of nucleotide processing rates and generation times that

Table 5. Growth, nutritional versatility and resource acquisition and storage: cellular characteristics that contribute to weed-like behaviour.^a

Metabolites, proteins, molecular characteristics, and other traits	Function(s)	Phenotypic traits	Notes and references
Ability to out-grow competitors Multiple copies of ribosomal DNA operons	Enhanced production of ribosomes	Rapid protein synthesis	Soil bacteria able to grow rapidly on high-nutrient media have an average of ~5.5 copies of ribosomal DNA operons (compared with ~1.4 copies for slow-growing species; Klappenbach <i>et al.</i> , 2000)
Key roles of di-/tripeptide-transport protein (<i>dtpT</i>) and oligopeptide transport ATP-binding protein (<i>Opp</i>) High-efficiency DNA polymerase	Permease (<i>dtpT</i>) and ABC-type transporter (<i>Opp</i>) that enable efficient scavenging of diverse peptides (Altermann <i>et al.</i> , 2005) Apparent correlation with short generation-time in fast-growing bacteria (Couturier and Rocha, 2006)	Efficient uptake of diverse peptides Rapid genome replication	These proteins enable species such as <i>Lactobacillus acidophilus</i> to efficiently scavenge peptides and thereby minimize the need for <i>de novo</i> synthesis of amino acids (Altermann <i>et al.</i> , 2005) The relatively fast-growing <i>Mycobacterium smegmatis</i> (generation time ~3 h) has a DNA polymerase processing rate of 600 nucleotide s ⁻¹ , compared with 50 nucleotide s ⁻¹ for <i>Mycobacterium tuberculosis</i> (generation time ~24 h; Straus and Wu, 1980; Couturier and Rocha, 2006)
Overlapping replication cycles	Bacteria can initiate a new round of chromosomal DNA replication before the completion of an ongoing round of synthesis	Rapid genome replication	<i>Escherichia coli</i> can operate up to eight origins of replication simultaneously (see Nordström and Dasgupta, 2006)
Polyploidy, aneuploidy, association with plasmids, and/or hybridization that enhance(s) vigour	Multiple copies of genomic DNA can lead to increased vitality, enhanced growth rates, increased production of antimicrobials, and enhanced energy generation and stress tolerance (see also Table 3)	Enhanced vigour	Shorter generation times in polyploid bacteria may arise from a number of factors including the increased number of ribosomal DNA operons (Cox, 2004; see also above) and gene localization within the genome (Pecoraro <i>et al.</i> , 2011). A recent study of F1 hybrids produced by crossing 16 wild-type strains of <i>S. cerevisiae</i> showed that many F1 strains displayed hybrid vigour (Timberlake <i>et al.</i> , 2011) ^b
Traits for high-efficiency energy generation (see Table 3) Efficient system to eliminate free radicals (see also Table 3)	Elimination of reactive oxygen species that can react with and damage macromolecular and cellular structures	Ability to cope with the oxidative stress associated with high growth rates	The enhanced rate of oxidative phosphorylation required for rapid growth result in the increased generation of free radicals; a phenomenon that has been observed in plant weeds (Wu <i>et al.</i> , 2011). <i>Aspergillus</i> mutants that lack superoxide dismutase genes are unable to grow under environmental conditions that favour a high metabolic rate (Lambou <i>et al.</i> , 2010). Strains of <i>Pseudomonas aeruginosa</i> have an auxiliary manganese-dependent version of superoxide dismutase that does not require iron and is upregulated under iron-limiting conditions (Britigan <i>et al.</i> , 2001)
Nutritional and metabolic versatility^c Genes for catabolism of a wide spectrum of hydrocarbons; e.g. those coding for the catechol-, homogenisate-, phenylacetate- and protocatechuate-catabolic pathways (<i>cat</i> , <i>hmg</i> , <i>pha</i> and <i>pca</i> gene families respectively) Catabolite-repression system for energy-efficient utilization of complex substrates Production and secretion of biosurfactants (e.g. rhamnolipids in <i>Pantoea</i> , <i>P. aeruginosa</i> and <i>P. putida</i> ; glycolipoproteins in <i>Aspergillus</i> spp.)	Facilitate the synthesis of proteins that enable the utilization of hydrocarbons (including polycyclic aromatic hydrocarbons) Advanced system to regulate utilization of diverse substrates Solubilization of hydrophobic molecules (and antimicrobial activity; see Table 6) Extracellular degradation of polymeric organic molecules for use as a carbon and energy source Oxidation of one-carbon substrates that are used for energy generation/growth Assimilation of nitrogen into anabolic pathways for <i>de novo</i> synthesis of amino acids and nucleic acids Glutamate dehydrogenase plays key roles in nitrogen assimilation and metabolism	Ability to utilize diverse hydrocarbons as a carbon and/or energy source Energy-efficient growth in nutritionally complex environments Ability to enhance the bioavailability of hydrophobic substrates Ability to grow saprotrophically Ability to utilize one-carbon substrates Ability to utilize a wide spectrum of nitrogen sources Ability to adapt metabolism for growth at high- or low-nutrient concentrations	Such compounds can be utilized as a sole carbon source by a small number of metabolically versatile microbes such as <i>Pantoea</i> spp., <i>Pseudomonas putida</i> and <i>Pichia anomala</i> (Timmins, 2002; Pan <i>et al.</i> , 2004; Vasileva-Tonkova and Gesheva, 2007) but are also known to exert cellular stress even at low concentrations (Bhaganna <i>et al.</i> , 2010). Catabolism of hydrocarbons will therefore alleviate stress at the same time as providing for energy generation and growth (Jiménez <i>et al.</i> , 2002; Domínguez-Cuevas <i>et al.</i> , 2006) A cAMP-independent catabolite-repression system has been identified in <i>P. putida</i> that enables the ordered assimilation of carbon, nitrogen and sulfur substrates (most energy-favourable first; Daniels <i>et al.</i> , 2010) <i>In vitro</i> studies of <i>Pantoea</i> and <i>P. aeruginosa</i> demonstrate that biosurfactants enhance the uptake of hydrophobic substrates (Al-Tahhan <i>et al.</i> , 2000; Vasileva-Tonkova and Gesheva, 2007; Reis <i>et al.</i> , 2011) and thereby facilitate colonization of hydrocarbon-containing habitats (see Table S1). <i>Aspergillus</i> species can produce copious amounts of biosurfactant (Desai and Banat, 1997; Kiran <i>et al.</i> , 2009) Saprotrophic capabilities enhance the range of substrates from which nutrition can be obtained. Saprotrophic enzymes from <i>Aspergillus</i> species can function over a wide range of environmental stresses (Hong <i>et al.</i> , 2001) One-carbon substrates are environmentally commonplace and their utilization can facilitate rapid growth (van der Klei <i>et al.</i> , 1991; McDonald and Murrell, 1997; Pacheco <i>et al.</i> , 2003) Versatility in the utilization of nitrogen substrates can enable proliferation in open habitats by, and opportunistic pathogenicity for, <i>Aspergillus</i> , <i>Pseudomonas putida</i> and other weed species (Kraupmann and Braus, 2005; Daniels <i>et al.</i> , 2010; see also Table S1). <i>Rhodotorula</i> spp., including metabolically versatile weed species, can grow on D-amino acids as a sole carbon and nitrogen source (Simoneita <i>et al.</i> , 1989; Pollegioni <i>et al.</i> , 2007) Expression of <i>gdh</i> was between 5- and 26-fold greater in <i>P. putida</i> cells under low-nutrient conditions compared with a high-nutrient control (Syn <i>et al.</i> , 2004). The synthesis of glutamate via glutamate dehydrogenase, rather than glutamate synthase, consumes 20% less ATP (Syn <i>et al.</i> , 2004)

Enzyme systems that enable metabolism under nutrient-limiting conditions by avoiding dependence on specific cofactors	Utilization of substitute cofactor, or replacement of enzyme system(s) that would require a cofactor that is environmentally scarce	Maintenance of essential enzyme functions at low nutrient concentrations	For example arginase in <i>S. cerevisiae</i> (Middelhoven <i>et al.</i> , 1969) and phospholipase C in <i>P. aeruginosa</i> (Stinson and Hayden, 1979)
Pathways to synthesize phosphate-free membrane lipids	Maintenance of membrane functions under phosphate-limited conditions	Ability to remain active, and grow, in phosphate-depleted environments	<i>Pseudomonas</i> spp. can synthesize membranous glycolipids as an alternative to phospholipids under low-phosphate conditions (Minnikin <i>et al.</i> , 1974). <i>Salinibacter ruber</i> and a number of phytoplankton species utilize sulfur in membrane glycolipids that have a sulfoquinovose-containing (non-phosphorus) head group and are synthesized in response to low phosphate availability (Benning, 1998; Corcelli <i>et al.</i> , 2004; Bellingier and Van Mooy, 2012)
High glucosylglycerate under low-nitrogen conditions	Control of nitrogen homeostasis to enable growth under nitrogen-limiting conditions	Nitrogen oligotrophy	Intracellular glucosylglycerate is increased in <i>M. smegmatis</i> under low nitrogen conditions. Mutants unable to synthesize glucosylglycerate showed reduced uptake of ammonium and a reduced growth rate (Behrends <i>et al.</i> , 2012)
Efficient acquisition and storage of resources			
Transport proteins to scavenge peptides and/or other macromolecules	Efficient scavenging of biomacromolecules avoiding the need for <i>de novo</i> synthesis	Efficient uptake of substrates	For example in <i>L. acidophilus</i> (see also above; Altermann <i>et al.</i> , 2005)
Biosurfactants to access hydrophobic substrates (see above)	Efficient degradation of complex extracellular substances	Efficient utilization of high- <i>M_r</i> substrates	For example, xylanases produced by <i>Acinetobacter junii</i> and <i>Aspergillus</i> spp. (Meagher <i>et al.</i> , 1988; Lo <i>et al.</i> , 2010)
Production of high quantities of high-affinity saprotrophic enzymes	Highly efficient uptake of diverse carbon substrates	Rapid uptake of hexose sugars	These proteins are upregulated in <i>S. cerevisiae</i> at low sugar concentrations (Wendell and Bisson, 1994)
High-affinity hexose-transport proteins, e.g. Hxt2p (<i>HXT</i> gene family)	Extracellular degradation of organic phosphate and uptake and intracellular polymerization of inorganic phosphate	Sequestration and storage of inorganic phosphate	<i>Pichia kudriavzevii</i> has a high rate of inorganic phosphate production from phytate (Helström <i>et al.</i> , 2012). Reduced survival of polyphosphate kinase-deficient bacterial mutants has been observed at low pH, high salinity, and under oxidative stress (Jahid <i>et al.</i> , 2006). Other weed species, such as <i>P. putida</i> , <i>Dunaliella salina</i> , <i>Microcystis aeruginosa</i> and <i>Ostreococcus tauri</i> are efficient accumulators of polyphosphate (Bental <i>et al.</i> , 1991; Shi <i>et al.</i> , 2003; Tobin <i>et al.</i> , 1997; 2007; Le Bihan <i>et al.</i> , 2011)
Phytase, inorganic phosphate transport proteins and polyphosphate kinase			The competitive ability of some bacterial weeds (e.g. pseudomonads, <i>Pantoea ananatis</i>) may be enhanced by siderophore production (Duff <i>et al.</i> , 1993; Loaces <i>et al.</i> , 2011). <i>Rhodotorula mucilaginosa</i> and <i>M. aeruginosa</i> upregulate siderophore production under iron-limiting conditions (Andersen <i>et al.</i> , 2003; Xing <i>et al.</i> , 2007); in contrast, <i>Saccharomyces</i> spp. are not known to produce siderophores (Lesuisse <i>et al.</i> , 2001)
Production and secretion of siderophores that chelate iron	To scavenge iron from the extracellular milieu which is then removed once the siderophores have been transported back into the cell	Efficient scavenging of iron	<i>Dunaliella salina</i> cells undergo a fourfold increase in iron-binding affinity under low-iron conditions. Iron is then taken up stored in vacuoles (Paz <i>et al.</i> , 2007)
Enhanced efficiency of iron-binding proteins (e.g. transferrin-like protein, TfT, in <i>D. salina</i>)	Binding and uptake of extracellular iron	Ability to accumulate iron under iron-limiting conditions	<i>Saccharomyces cerevisiae</i> stores arginine (in vacuoles) that can be utilized when the extracellular environment becomes nitrogen substrate limited (Westenberg <i>et al.</i> , 1989)
Arginine biosynthesis proteins	Bulk synthesis of arginine as a nitrogen substrate store	Amino acid accumulation and storage	Trehalose synthesis can be upregulated prior to the stationary growth phase and during stress (Parrou <i>et al.</i> , 1997), and glycogen synthesis is upregulated during periods of nutrient limitation, in <i>S. cerevisiae</i> (Lillie and Pringle, 1980; Farkas <i>et al.</i> , 1990). Starch is accumulated by <i>D. salina</i> (Stoyanova-Bakalova and Toncheva-Panova, 2003)
Trehalose-synthesis proteins (e.g. TPS, TPP) and glycogen synthase (e.g. GSY1 and GSY2)	Synthesis of disaccharide- and polysaccharide-storage compounds (e.g. trehalose and glycogen respectively)	Effective synthesis and storage of carbohydrates	Polyhydroxybutyrate, in the form of intracellular granules, dominates the cytoplasm of <i>Halobacterium salinarum</i> (Saponetti <i>et al.</i> , 2011); <i>Dunaliella</i> spp. also store lipids (Chen <i>et al.</i> , 2011); recent studies of <i>P. putida</i> suggest that carbon and energy can be channelled into storage of polyhydroxybutyrate without a reduction of growth rate (Escapa <i>et al.</i> , 2012)
Polyhydroxybutyrate-synthesis proteins (and those producing other storage lipids)	Synthesis of lipid stores (see also Table 3)	Bulk storage of lipids	The remarkable morphology of <i>H. walsbyi</i> cells optimizes acquisition of resources in highly competitive high-salt habitats (Bohujs <i>et al.</i> , 2006; Bardavid <i>et al.</i> , 2008)
Morphological and physiological adaptations to optimize light capture and absorption of oxygen and nutrients	Flat cell shape gives a high surface-to-volume ratio; gas vesicles help to maintain the horizontal orientation of planktonic cells	Optimized light capture, and oxygen and nutrient absorption	Habitat dominance by <i>Gonyostomum semen</i> can be attributed, in part, to having a higher photosynthetic rate than other autotrophic species (Coleman and Heywood, 1981; Peltomaa and Ojala, 2010)
Efficient chlorophyll <i>a</i> and chloroplast arrangement	Optimization of light harvesting and high photosynthetic rate	High photosynthetic rate	Twelve genes involved in gas synthesis and storage have been identified in <i>M. aeruginosa</i> (Mlouka <i>et al.</i> , 2004); models suggest regulation of buoyancy in <i>M. aeruginosa</i> has a key role in habitat dominance (Bonnet and Poulin, 2002)
Genes related to gas synthesis and gas vesicle formation (e.g. <i>gvp</i> family in <i>M. aeruginosa</i>)	Gas vesicles provide buoyancy to optimize access to light and high oxygen concentrations (close to the water:atmosphere interface of aquatic habitats)	Ability to access resources required for photosynthesis	

a. The current table provides examples of proteins, genes or other characteristics that can be associated with and give rise to weediness; the categories of traits are not mutually exclusive. Individual traits may not be unique to microbial weeds; however, weeds species are likely to have a number of these types of characteristics.

b. Whereas polyploidy can enhance vigour (via increased growth rates, enhanced stress tolerance etc.) increased chromosome number can also result in decreased growth rates; the consequences of polyploidy can vary with species and environmental context.

c. Nutritional versatility is particularly advantageous for microbial weed species that have a generalist lifestyle such as *Pseudomonas putida* and *Pichia anomala* (Timmis, 2002; Walker, 2011).

are an order of magnitude faster than those of comparable *Mycobacterium* species (Table 5). Light and/or oxygen are key resources for many planktonic species; 12 genes have been identified that are involved in gas synthesis and storage in species such as *M. aeruginosa* and *H. walsbyi* (Table 5). Modelling approaches indicate that buoyancy, which enables the cell to access the upper (well-illuminated, well-oxygenated) section of the water column, plays a key role in habitat dominance by *M. aeruginosa* (Bonnet and Poulin, 2002). Studies of *H. walsbyi* suggest that its peculiar, flat cell-shape optimizes the uptake of nutrients and oxygen and its intracellular gas vesicles help to maintain a horizontal orientation to capture light (Table 5). In freshwater habitats dominance of algal blooms by *G. semen* has been attributed in part to a high photosynthetic rate that results from a combination of its high-efficiency chlorophyll *a* and a highly efficient chloroplast arrangement (Table 5).

In open habitats that are nutritionally dynamic and/or complex (Table S1) metabolic versatility in nutrient utilization can enhance the competitive ability of microbes (for examples see Table 5). *Pseudomonas putida* and related species are remarkable in their range of nutritional strategies (Table 5; Timmis, 2002; 2009): they are able to catabolize an indeterminate number of hydrocarbons (e.g. decanoate, toluene) and other organic molecules (e.g. amino acids, and organic acids such as acetic, citric and lactic acid) for use as carbon and energy sources; amino acids, dipeptides, ammonium chloride, urea, ammonium nitrate, and other substances as nitrogen sources; amino acids, organic acids, sodium sulfite, thiouracil, agmatine sulfate, etc. as sulfur sources; as well as accumulating diverse substances as intracellular reserves (Chakrabarty and Roy, 1964; Daniels *et al.*, 2010; Table 5) enabling a switch to oligotrophic growth at low nutrient concentration (Table 5; Timmis, 2002). In addition, *P. putida* and closely related species have evolved a complex, cAMP-independent catabolite repression system which incorporates five different signals and enables an ordered assimilation of carbon, nitrogen and sulfur substrates from the most to least energetically favourable (Daniels *et al.*, 2010). This may contribute to the ability of many pseudomonads to sustain vigorous growth in nutritionally complex habitats. Pseudomonads, and closely related bacteria such as *Acinetobacter*, are not only environmentally ubiquitous but can dominate microbial communities in numerous types of open habitat including the plant phyllosphere and rhizosphere, bioaerosols, oil sands, and other hydrocarbon-containing environments (see Table S1; Ogino *et al.*, 2001; Röling *et al.*, 2004; Louvel *et al.*, 2011). *Pseudomonas* species are able to enhance the availability of hydrophobic substrates by secreting biosurfactants and, at the same time, tolerate the stress induced by the latter (see below). Despite their copio-

trophic phenotype, pseudomonads utilize a number of strategies to maintain growth and metabolism under phosphate-, iron- or nitrogen-depleted conditions (Table 5), can synthesize glutamate via glutamate dehydrogenase rather than glutamate synthase (thereby consuming 20% less ATP; Syn *et al.*, 2004), and have even been observed growing in distilled water (Favero *et al.*, 1971; Hirsch, 1986). Specific enzymes of *Pseudomonas aeruginosa* are catalytically active even when cofactor nutrients are scarce, and some strains have a manganese-dependent superoxide dismutase which does not require iron and is upregulated at low iron concentrations (Table 5).

Weeds such as lactic acid bacteria and *S. cerevisiae* that lack saprotrophic capabilities can dominate communities in habitats with high nutrient concentrations (Tables S1 and 5); they possess high-affinity transport proteins that enable rapid and efficient uptake of substrate molecules to support their vigorous growth phenotypes (see Table 5). Cells of *S. cerevisiae* are not only highly efficient at storing carbohydrates (combined trehalose and glycogen may reach 25% dry weight; Aranda *et al.*, 2004; Guillou *et al.*, 2004), but they can also accumulate high concentrations of arginine and polyphosphate and have adaptations to retain activity under low-nitrogen conditions (Table 5). Plant weeds that excel in open habitats have distinct genetic properties; they are often of hybrid origin and/or polyploid, traits that give rise to increased growth rates and stress tolerance, and have specific genes that confer their vigorous phenotype (Anderson, 1952; Hill, 1977; Schnitzler and Bailey, 2008; Bakker *et al.*, 2009). There is also evidence that the multiple copies of genomic material found in vigorous microbes that enhanced vigour have arisen via hybridization and/or increased ploidy levels in bacterial weed species, as well as *Pichia* and *Saccharomyces* (Table 5; Cox, 2004). A recent study by Timberlake and colleagues (2011) reported vigorous growth for the majority of F1 hybrids produced by crossing 16 wild-type *S. cerevisiae* strains (although hybridization and/or polyploidy may not always be advantageous; see Table 5). Many microbial weed species are phenotypically variable (like their plant counterparts; Table 1; Hill, 1977) and may form part of a species complex due to polyploidization and hybridization (e.g. *Acinetobacter*, *Pseudomonas*, *Mycobacterium*, *Saccharomyces sensu stricto*; Table S1; Rainieri *et al.*, 1999; Todd *et al.*, 2004; Ballerstedt *et al.*, 2007; Parkinson *et al.*, 2011; Sicard and Legras, 2011); such factors may contribute to the vigour and weediness of such species.

High- M_r macromolecules are the primary microbial substrates available in many habitats: e.g. hydrocarbons (in bioaerosols, oil-containing environments, and the rhizosphere), and polysaccharides (within or derived from plant necromass such as cellulose, lignin and humic

substances; see Table S1). In order to out-compete other species in such habitats, microbes must utilize these macromolecular substrates that typically have a low bioavailability, require specialist enzymes to degrade, can be toxic, and/or are energy-expensive to catabolize. Facultatively saprotrophic microbes such as *Aspergillus*, *Pichia*, *Rhodotorula* and *Acinetobacter* species are able to access and catabolize recalcitrant polysaccharides; some weed species can produce exceptional amounts, or high-affinity versions, of inducible saprotrophic enzymes, such as xylanases (Table 5). Several saprotrophic enzymes produced by *Aspergillus* species (that can dominate physicochemically diverse habitats; Table S1) can retain activity over a wide range of environmental stresses (Hong *et al.*, 2001); and *Aspergillus*, and *Rhodotorula* spp. are able to utilize a broad range of nitrogen substrates thereby avoiding the need for *de novo* synthesis (Table 5). Whereas nutritionally complex habitats can select for metabolic versatility (see below) and thereby favour the growth of a great number of generalist species, only a minute fraction of such microbes are able to dominate what were (initially) phylogenetically diverse microbial communities (Table S1).

Antimicrobial strategies and trophic interactions

Grazing, predation and viral infections adversely impact the growth dynamics of and/or incur massive losses on microbial populations and so exert a powerful selective pressure, and this can be well illustrated by the ecologies of many planktonic species in aquatic habitats (Matz *et al.*, 2004; Jüttner *et al.*, 2010; Berdjeb *et al.*, 2011; Thomas *et al.*, 2011; Yang and Kong, 2012). The competitive ability of some aquatic bacteria and eukaryotes (both marine and freshwater species) is associated with their ability to resist or tolerate virus infection (see Table 6). Species with a halophilic growth phenotype, especially those which have high growth rates in salt-saturated environments such as *S. ruber* and *H. walsbyi* minimize losses to predators because these salt concentrations are hostile to protozoa due to their low water activity, high ionic strength, and/or (for high-MgCl₂ and high-CaCl₂ brines) high chaotropicity (Table 6; Hallsworth *et al.*, 2007). *Gonyostomum semen* has evolved vertical migration patterns through the water column that are out of phase with those of grazing species and thereby minimizes losses (Table 6). Other weeds, including *P. aeruginosa* and *M. aeruginosa*, are capable of forming multicellular structures – microcolonies – that are sufficiently large to reduce losses by many species of grazers and predators (Table 6). Species such as *M. aeruginosa* accumulate toxic metabolites, and *M. aeruginosa* and *Aspergillus* spp. are known to synthesize secondary metabolites that repel to grazing plankton and arthropods respectively (Table 6).

A number of agricultural (plant) weeds are able to dominate their habitats due largely to the production of phytotoxic inhibitors that can be exuded from leaves, roots and plant necromass and which inhibit the germination and/or growth of other plant species (Hill, 1977; Xuan *et al.*, 2006; Alsaadawi and Salih, 2009; Chou, 2010; Meksawat and Pornprom, 2010; Inderjit *et al.*, 2011). It is likely that most microbial weed species also secure their competitive advantage – in part at least – via the deployment of antimicrobial substances. Assuming that the examples listed in Table S1 are representative of the microbial biosphere then the proportion of habitats dominated by eukaryotic species is approximately 40%. If a short generation time was sufficient to facilitate habitat domination then prokaryotic species would populate almost all these habitats. The fact that species of algae, fungi and yeast are also able to do so underscores the importance of factors other than generation time in the biology of habitat dominance. *Saccharomyces cerevisiae* provides an illustration of how one weed species combines the synthesis of a potent antimicrobial, energy generation, and coordinated synthesis of stress protectants in a single process, i.e. aerobic production of the chaotropic alcohol, ethanol (see above; Table 6). Ethanol can be produced and secreted at concentrations > 4 M that cause a reduction of water activity (to 0.896) and exert an exceptionally high chaotropic activity (see above) sufficient to prevent the growth or metabolism of any living system (Hallsworth, 1998; Hallsworth and Nomura, 1999; Hallsworth *et al.*, 2007); *S. cerevisiae* simultaneously produces glycerol (which can reduce ethanol stress in yeast and fungi; Hallsworth, 1998; Hallsworth *et al.*, 2003b) during ethanol fermentation and is metabolically wired to synthesize high concentrations of the protectant trehalose at the end of its growth cycle when ethanol concentration is highest (see Table 6).

We believe that the bulk production and secretion of chaotropic solutes and organic acids (that can reach molar concentrations) is unique to microbial weed species (Tables 1 and 6). Whereas chaotropic substances can promote growth and expand the biotic windows of microbes under specific conditions (Hallsworth and Magan, 1994b; 1995; Williams and Hallsworth, 2009; Bhaganna *et al.*, 2010; Chin *et al.*, 2010), in other circumstances chaotropicity can be highly inhibitory or even lethal (Hallsworth, 1998; Hallsworth *et al.*, 2003a; 2007; Duda *et al.*, 2004; Bhaganna *et al.*, 2010). For plant weeds it is the early stages of habitat colonization that are key to their success (Hill, 1977), and it may be that the efficient synthesis of antimicrobials is frequently as important as much as the type of substance produced (Hallsworth, 1998; Pretorius, 2000; Eshkol *et al.*, 2009; Fialho *et al.*, 2011). Unlike plant weeds, where phytotoxic inhibitors tend to have the greatest impact on neighbouring plants, microbes can also produce volatile

Table 6. Antimicrobial activities and trophic interactions: cellular characteristics that contribute to weed-like behaviour.^a

Metabolites, proteins, molecular characteristics, and other traits	Function(s)	Phenotypic traits	Notes and references
Avoidance of/resistance to viral infection, predators, and grazers Metabolites that repel grazers and/or predators, e.g. aldehydes and other volatile organic compounds in plankton and fungi	Secondary metabolites produced by <i>Aspergillus</i> spp. repel (and can reduce the fecundity of) grazing arthropods such as <i>Folsomia candida</i> ; those produced by <i>Microcystis aeruginosa</i> act as a repellent to planktonic grazers	Ability to repel grazers and predators	<i>Aspergillus nidulans</i> mutants lacking the <i>laeA</i> gene, a global regulator of secondary metabolite synthesis, are consumed by <i>F. candida</i> in preference to the wild-type strain (Rohlfis et al., 2007). Aldehydes produced by plankton repel crustacean grazers (Jüttner, 2005); β -cyclocitral from <i>M. aeruginosa</i> has been shown to repel <i>Daphnia magna</i> (Jüttner et al., 2010)
Synthesis of extracellular polymeric substances (e.g. <i>M. aeruginosa</i>), or use of flagella (e.g. <i>Pseudomonas aeruginosa</i>), that facilitate microcolony formation	Formation of multicellular structures that are sufficiently large to minimize grazing and predation by single-celled protozoa, zooplankton etc.	Microcolony formation	The role of polymeric substances in colony formation has been established using <i>M. aeruginosa</i> as a model organism (Yang and Kong, 2012). Functional analysis of <i>P. aeruginosa</i> mutants that were unable to form flagella, or were flagellate-deficient, demonstrated an inability to form microcolonies that can confer resistance to grazing (Matz et al., 2004)
Biochemical adaptations to resist viral infection	Diverse mechanisms including modification of cell surface to prevent binding, intracellular suppression or viral genome replication, and degradation of viral DNA by restriction endonucleases	Resistance to or tolerance of viral infection	There are a number of mechanisms by which dominating species of bacteria and eukaryotic plankton (e.g. <i>P. aeruginosa</i> , <i>Ostreococcus tauri</i> ; <i>Bathycoccus</i> and <i>Micromonas</i> spp.) may acquire and maintain resistance to viral infection (see Thomas et al., 2011; Selezska et al., 2012)
Toxic metabolites (e.g. microcystin in <i>M. aeruginosa</i>)	Toxicity against grazing or predatory zooplankton	Ability to protect against excessive grazing	Microcystin synthesis in <i>M. aeruginosa</i> can be induced by exposure to zooplankton; concentrations of the toxin in cells exposed to <i>D. magna</i> , <i>Daphnia pulex</i> and <i>Moina macrocarpa</i> were up to fivefold those found in control cells (Jiang et al., 2003)
Halophilic growth phenotype	Colonization of salt-saturated habitats that are hostile to predatory protozoa and minimizes losses from grazing and predation	Ability to inhabit environments hostile to predators and grazers	Halophiles that dominate high-salt habitats (e.g. <i>D. salina</i> , <i>S. ruber</i>) minimize interactions with grazers and predators most of which cannot tolerate molar salt concentrations (Bardavid et al., 2008)
Biochemical and cellular infrastructure that give rise to idiosyncratic migration patterns in the water column	Vertical migration within the water column that minimizes loss through predation and grazing	Avoidance of predators and grazers	The predominance of <i>Gonyostomum semen</i> in eutrophic habitats has been associated in part with migration patterns that minimize loss through grazing (Salonen and Rosenberg, 2000)
Bulk production of highly soluble, moderately chaotropic stressors^b Ethanol produced via a high- K_m alcohol dehydrogenase (e.g. Adh1 in <i>Saccharomyces cerevisiae</i>) Enzymes for acetone, isopropanol and butanol synthesis	Efficient conversion of acetaldehyde to ethanol; in contrast, low- K_m Adh typically converts ethanol to acetaldehyde under <i>in vivo</i> conditions (Piskur et al., 2006) Synthesis of chaotropic substances in the anaerobic bacterium <i>Clostridium acetobutylicum</i>	Production and secretion of ethanol (up to molar concentrations) Bulk production, and secretion, of multiple chaotropic stressors	Ethanol, secreted in quantity by species such as <i>S. cerevisiae</i> and <i>Zymomonas mobilis</i> , is a moderately chaotropic substance that is nevertheless a potent stressor at millimolar to molar concentrations (Hallsworth, 1998; Bhaganna et al., 2010; Cray et al., 2013) Acetone, isopropanol and butanol that are produced by <i>C. acetobutylicum</i> (George et al., 1983) are more chaotropic ethanol – when compared on a molar basis – and are therefore more potent in their inhibitory activity as cellular stressors (Bhaganna et al., 2010; Cray et al., 2013; Bell et al., 2013)
Production of high concentrations of urea and ammonia	Synthesis of low- M_r , nitrogenous chaotropes	Production and secretion of urea and ammonia at inhibitory concentrations	Urea and ammonia are produced by phylogenetically diverse bacteria such as <i>Pseudomonas stutzeri</i> and <i>Clostridium aminophilum</i> respectively (Therkildsen et al., 1997; Watanabe et al., 1998; Anderson et al., 2010)
Release of VOCs as low-solubility antimicrobial stressors Highly volatile metabolites that act as hydrophobic stressors (log $P > 1.95$) ^c	Inhibition of microbial metabolism; typically via hydrocarbon-induced water stress (Bhaganna et al., 2010) Inhibition of microbial metabolism via chaotrope-induced water stress (Hallsworth, 1998; Hallsworth et al., 2003a; Bhaganna et al., 2010)	Production and release of volatile antimicrobials Production and release of volatile antimicrobials	Examples are: isoamyl acetate (produced by <i>Pichia anomala</i> ; Passoth et al., 2006), ethyl octanoate (<i>S. cerevisiae</i> ; Fialho et al., 2011), β -cyclocitral (<i>M. aeruginosa</i> and <i>Dunaliella salina</i> ; Herrero et al., 2006; Ozaki et al., 2008), pentane (<i>D. salina</i> ; Muñoz et al., 2004) and diverse substances from <i>Aspergillus</i> (see above), <i>R. mucilaginosa</i> (Bazzini et al., 2005), <i>P. putida</i> and other <i>Pseudomonas</i> species (Morales et al., 2005) Examples are: ethyl acetate (produced by <i>Pichia anomala</i> ; Walker, 2011), dichloromethane (<i>Dunaliella</i> spp.; Colomb et al., 2008) and 2-phenylethanol (<i>Mycobacterium bovis</i> , <i>Mycobacterium smegmatis</i> and <i>S. cerevisiae</i> ; Eshkol et al., 2009; McNamey et al., 2012); and a range of chaotropic stressors by <i>R. mucilaginosa</i> (Bazzini et al., 2005), <i>P. putida</i> and other <i>Pseudomonas</i> species (Morales et al., 2005)

<p>Volatile catabolites produced from the degradation of macromolecular substrates</p> <p>Acidification of the extracellular environment Secretion of organic acids</p> <p>High-activity L-lactate dehydrogenase (e.g. <i>ldhL</i> in <i>Lactobacillus plantarum</i>)</p> <p>Enzymes for acetic and butyric acid synthesis (e.g. phosphotransacetylase and acetate kinase, and phosphate butyryltransferase and butyrate kinase respectively)</p> <p>Production of biosurfactants, enzymes, and/or toxins with antimicrobial activities Biosurfactants (e.g. rhamnolipids in <i>Pseudomonas</i> spp.; glycolipoproteins in <i>Aspergillus</i> spp.)</p> <p>Panomycolin (an α-D-glucanase)</p> <p>Ochratoxin and other mycotoxins (e.g. in <i>Aspergillus</i> spp.)</p> <p>Antifungal peptide (Anafp, in <i>Aspergillus niger</i>)</p> <p>Mycocins (e.g. salt-mediated killer toxin)</p> <p>Bacteriocins e.g. plantaricins (<i>plh</i> gene family), lactacin-B (La1791–La1803), plant lectin-like bacteriocin (LlpA)</p> <p>Pyoluteorin (in <i>P. putida</i>)</p> <p>Phenazine and phenazine-derived antimicrobials</p> <p>Mechanisms to induce cell lysis</p> <p>Resistance to antimicrobial stressors and toxins Coordinated production of compatible solutes and ethanol</p>	<p>Inhibition of microbial metabolism typically via chaotrope-induced or hydrocarbon-induced water stress (Hallsworth <i>et al.</i>, 2003a; Bhagana <i>et al.</i>, 2010)</p> <p>Acidification of the extracellular milieu</p> <p>Intracellular conversion of pyruvate to lactate</p> <p>Conversion of acetyl-CoA to acetate; and butyryl-CoA to butyrate</p> <p>Production of biosurfactants, enzymes, and/or toxins with antimicrobial activities Solubilization of hydrophobic substrates; inhibitory to the growth of many microbial competitors</p> <p>Hydrolysis of glucans such as laminarin</p> <p>Inhibition of other microbial species/strains</p> <p>Mode-of-action not yet resolved</p> <p>Inhibitory to many fungi, yeasts and other microbes</p> <p>Modification of membrane structure causing leakage and dissipation of proton-motive force (Andersen <i>et al.</i>, 1998; Parret <i>et al.</i>, 2003; Altermann <i>et al.</i>, 2005)</p> <p>Inhibitory to growth of oomycetes and fungi; mode-of-action not yet resolved</p> <p>Probable antimicrobial activity</p> <p>Elimination of potential competitors</p> <p>Protection against ethanol stress</p>	<p>Production and release of volatile antimicrobials⁹</p> <p>Ability to acidify the habitat</p> <p>Production and secretion of lactic acid</p> <p>Production/secretion of acetic acid and butyric acid</p> <p>Production of inhibitory biosurfactants</p> <p>Antimicrobial and cytocidal activity via degradation of cell wall components</p> <p>Secretion of antimicrobials</p> <p>Secretion of an antifungal peptide</p> <p>Production of antibiotic peptides</p> <p>Production of antibacterial peptides</p> <p>Antifungal activity</p> <p>Antimicrobial activity</p> <p>Algicidal activity</p> <p>Resistance to ethanol</p>	<p>Antimicrobial stressors can be released during the saprotrophic degradation of lignin, catabolism of hydrocarbons etc. (e.g. catechol, vanillin and biphenyl in <i>P. putida</i>; Zuroff and Curtis, 2012)</p> <p><i>Acinetobacter rhizosphaerae</i> and <i>S. cerevisiae</i> secrete a range of substances that can acidify the environment (Kotyk <i>et al.</i>, 1999; Gulati <i>et al.</i>, 2010); <i>Aspergillus</i> spp. produces high levels of citric acid (up to 200 g l⁻¹ in 5–7 days; Ward <i>et al.</i>, 2006)</p> <p>The specific activity of <i>ldhL</i> from a lactic acid produced by <i>L. plantarum</i> was found to be 2460 U mg⁻¹; compared with 250 U mg⁻¹ for <i>Escherichia coli</i> (Garvie, 1980). Habitat acidification by lactic acid inhibits the growth of many microbial competitors (Table S1; Guillot <i>et al.</i>, 2003). <i>Lactobacillus</i> efficiently acidifies its environment even at low sugar concentrations (Klinke <i>et al.</i>, 2009)</p> <p><i>Clostridium</i> spp. produce acetic, butyric and lactic acids; fermentations carried out in media with an initial pH of ~6.5 resulted in a final pH of 4–4.5 (Wiesenborn <i>et al.</i>, 1989; Boynton <i>et al.</i>, 1996; Liu <i>et al.</i>, 2011)</p> <p>Lipopeptides from <i>Pseudomonas</i> sp. DF 41 were shown to inhibit <i>Sclerotinia sclerotiorum</i> (Berry <i>et al.</i>, 2010); purified biosurfactants from <i>Pseudomonas fluorescens</i> inhibit growth of <i>Botrytis cinerea</i> and <i>Rhizoctonia solani</i> at low concentrations (25 µg ml⁻¹) and can lyse zoospores of <i>Phytophthora capsici</i> (Kruit <i>et al.</i>, 2009). Studies of <i>P. aeruginosa</i> rhamnolipid suggests a key role in competitive ability against other bacteria (Wecke <i>et al.</i>, 2011). <i>Acinetobacter</i> spp. produces a potent biosurfactant (phosphatidylethanolamine; Desai and Banat, 1997) which may act as an antimicrobial. Biosurfactants from <i>Aspergillus ustus</i> are inhibitory to phylogenetically diverse microbes (Kiran <i>et al.</i>, 2009)</p> <p>Panomycolin is produced by <i>P. anomala</i> and is inhibitory or lethal to many species of yeasts and fungi such as <i>B. cinerea</i> (Izgü <i>et al.</i>, 2005)</p> <p>Mycotoxins can inhibit phylogenetically diverse microbes (including fungal and bacterial species) and thereby enhance competitive ability (Boultonnes <i>et al.</i>, 1984; Magan <i>et al.</i>, 2003)</p> <p>Anafp inhibits growth of yeasts and fungi including some <i>Aspergillus</i> and <i>Fusarium</i> spp., <i>Candida albicans</i>, <i>S. cerevisiae</i> and <i>Trichosporon beigelii</i> (Lee <i>et al.</i>, 1999)</p> <p>A range of killer toxins produced by <i>Pichia farinosa</i>, <i>Rhodotorula</i> spp. and <i>S. cerevisiae</i> that are inhibitory to phylogenetically diverse taxa (Golubev and Churkina, 1993; Golubev <i>et al.</i>, 1996; Suzuki <i>et al.</i>, 2001; Trindade <i>et al.</i>, 2002; Abergaria <i>et al.</i>, 2010)</p> <p>For example <i>L. plantarum</i>, <i>Lactobacillus acidophilus</i> and <i>Pseudomonas</i> spp.</p> <p>Pyoluteorin-producing <i>P. putida</i> strains have potential as biocontrol agents of pathogens such as <i>Glomerella lucumanensis</i> (sugarcane red rot; Nowak-Thompson <i>et al.</i>, 1999; Hassan <i>et al.</i>, 2011)</p> <p>Produced by species such as <i>M. aeruginosa</i> and <i>P. aeruginosa</i> (Mavrodi <i>et al.</i>, 2006; Lifshits and Carmeli, 2012)</p> <p><i>Gonyosomum semen</i> is able to lyse microbial cells (e.g. <i>Rhodomonas lacustris</i>) on contact; mechanism not yet resolved (Rengefors <i>et al.</i>, 2008)</p> <p>The synthesis of glycerol (known to afford some protection against ethanol stress; Hallsworth, 1998; Hallsworth <i>et al.</i>, 2003b) and ethanol production are coupled in <i>S. cerevisiae</i>; accumulation of trehalose (an effective protectant against ethanol; Mansure <i>et al.</i>, 1994) coincides with the growth phase when ethanol concentrations are highest</p>
--	---	--	--

Table 6. cont.

Metabolites, proteins, molecular characteristics, and other traits	Function(s)	Phenotypic traits	Notes and references
Compatible-solute accumulation in response to solvent and hydrophobic stressors (see above)	Protection of macromolecular systems challenged by chaotropic and hydrophobic stressors (see studies of <i>A. nidulans</i> , <i>S. cerevisiae</i> and <i>P. putida</i> ; Mansure et al., 1994; Hallsworth et al., 2003b; Park et al., 2007; Bhaganna et al., 2010; Bell et al., 2013)	Ability to maintain functionality of macromolecular systems	Kosmotropic compatible-solutes and other kosmotropic substances can reduce stress induced by chaotropic solutes such as MgCl ₂ , glycerol and ethanol (Hallsworth, 1998; Hallsworth et al., 2007; Williams and Hallsworth, 2009). Tryptophan and trehalose are upregulated in cells of both eukaryotic and bacterial species in response to ethanol (Parrou et al., 1997; Purvis et al., 2005; Hirasawa et al., 2007; Horinouchi et al., 2010). <i>S. cerevisiae</i> could tolerate 28% v/v ethanol when protected by the kosmotropic compatible-solute proline (Thomas et al., 1993; Hallsworth, 1998)
Upregulation of protein-stabilization proteins (see Table 3)	Protection of macromolecular systems from perturbations induced by antimicrobial stressors	Maintenance of protein structure and function during stressor-induced challenges	<i>Saccharomyces cerevisiae</i> upregulates synthesis of up to 10 heat-shock proteins under ethanol stress (Ma and Liu, 2010). <i>Pseudomonas putida</i> synthesizes diverse protein-stabilization proteins in response to chaotropic and hydrophobic stressors (see above; Hallsworth et al., 2003a; Bhaganna et al., 2010)
Membrane-stabilization proteins; solvent pumps (see Table 3) Multidrug efflux pumps (e.g. AcrAB-TolC in <i>E. coli</i> ; Pdr5p in <i>S. cerevisiae</i> ; ABC transporter proteins in <i>Aspergillus</i> spp.)	Expulsion of chemically diverse stressors and toxins	Tolerance to a wide array of stressors and toxins	Including chaotropic solutes (e.g. phenol, benzyl alcohol), hydrophobic stressors (e.g. <i>n</i> -hexane, <i>p</i> -xylene) and antibiotics (e.g. chloramphenicol, tetracycline, novobiocin) by AcrAB-TolC; fluorescent dyes (e.g. daunorubicin, rhodamine), drugs (e.g. tamoxifen); and ionophoric peptides (e.g. nigericin) by Pdr5p (Kolaczowski et al., 1996); and antifungals (clotrimazole and echinocandin B) by the <i>Atf1MDR1</i> gene product (Tobin et al., 1997; Nishino and Yamaguchi, 2001; Ramos et al., 2002)
Proteins for expulsion of, or resistance to, bacteriocins; e.g. lincomycin-resistance protein (lincR), undercrapenyl pyrophosphate phosphatase (bacA)	Removal of bacteriocins from the cell	Resistance to bacteriocins	LmrB is an ATP-binding cassette-type transporter in <i>Lactococcus lactis</i> (Gajic et al., 2003); BacA is a membrane-associated protein that in <i>E. coli</i> for (El Ghachi et al., 2004)
Enzymes that inactivate antimicrobial substances	Detoxification and/or catabolism of inhibitory stressors and toxins	Ability to detoxify or degrade inhibitory antimicrobials	Examples include degradation of β -lactam antibiotics in <i>Achromobacter</i> spp. and <i>P. aeruginosa</i> (Langhaee et al., 2000; Sinha et al., 2007) and degradation of volatile stressors such as dichloromethane (e.g. <i>Bacillus</i> spp.; Wu et al., 2009)
Enhanced tolerance to a chaotropic volatile	Resistance to 2-phenylethanol	Resistance to VOCs	One strain of <i>S. cerevisiae</i> was found to be capable of growth at up to ~24 mM 2-phenylethanol (Eshkol et al., 2009). For a chaotropic stressor, log P 1.52, almost as chaotropic as phenol, this is an exceptional level of tolerance (see Bhaganna et al., 2010)
Cell wall composition that confers resistance to specific antibiotics	Pathways targeted by specific antibiotics may be absent; cell wall can act as a barrier to antimicrobials	Resistance to antibiotics	The <i>Halobacterium walsbyi</i> cell wall is not composed of peptidoglycan so this species has reduced sensitivity to β -lactam antibiotics (Bardavid and Oren, 2008); the <i>M. smegmatis</i> cell envelope contains mycolic acids that reduce permeability, and therefore enhance resistance, to stressors and antibiotics (Liu and Nikaido, 1999)
Synthesis of extracellular polymeric substances (see Table 3)	Production and secretion of polymeric substances	Resistance to diverse antimicrobials	For example production of alginate by <i>P. aeruginosa</i> increases resistance to antibiotics such as tobramycin (Hentzer et al., 2001)

a. The current table provides examples of proteins, genes or other characteristics that can be associated with and give rise to weediness; the categories of traits are not mutually exclusive. Individual traits may not be unique to microbial weeds; however, weeds species are likely to have a number of these types of characteristics.

b. Chaotropic activity of > 5 and < 40 kJ kg⁻¹ mole⁻¹ (Cray et al., 2013).

c. Volatile organic compounds are unlikely to exceed micromolar concentrations in the cytosol of the weed cell.

d. Glycerol is a phylogenetically ubiquitous as a compatible solute for eukaryotic microbes, yet at molar concentrations it acts as a chaotropic stressor (Williams and Hallsworth, 2009; Cray et al., 2013).

organic compounds (VOCs) that inhibit the growth of both proximal and distal cells via their dispersion in both aqueous and gaseous milieu (Table 6).

Volatile organic compounds – including aldehydes, furones, terpenes and many alcohols – can be powerful inhibitors of microbial metabolism and growth (Table 6; Jorge and Livingston, 1999; Passoth *et al.*, 2006; Minerdi *et al.*, 2009; Fialho *et al.*, 2011; Corcuff *et al.*, 2011). Despite their volatility they are to some degree also water-soluble, and are sufficiently inhibitory to prevent metabolic activity and growth, even in aqueous solution at subsaturated solute-concentrations (Bhaganna *et al.*, 2010). VOCs with a $\log P_{\text{octanol-water}} < 1.95$ typically partition into the aqueous phase of cellular macromolecules and can act as potent chaotropic stressors (Table 6; Hallsworth *et al.*, 2003a; Bhaganna *et al.*, 2010; Cray *et al.*, 2013). Those stressors with a $\log P$ of > 1.95 preferentially partition into the hydrophobic domains of membranes and other macromolecular systems, and thereby act as hydrophobic stressors that induce a distinct type of chaotropicity-mediated water stress (Bhaganna *et al.*, 2010). We believe that the primary ecophysiological value of VOCs for microbial cells pertains to competition, species succession and (for weed species) habitat dominance. *Saccharomyces cerevisiae* synthesizes both hydrophobic and chaotropic volatiles that have antimicrobial activities, including ethyl octanoate and isoamyl acetate, and ethyl acetate and 2-phenylethanol (respectively, see Table 6; Fialho *et al.*, 2011). Ethanol and 2-phenylethanol are synergistic in their antimicrobial activity (Ingram and Buttke, 1984; Seward *et al.*, 1996), and this is consistent with their common chaotropicity-mediated mode-of-action. The more potent chaotrope of the two (when compared on a molar basis) is 2-phenylethanol which prevents growth of microbes in the range 10–40 mM (Berrah and Konetzka, 1962; Eshkol *et al.*, 2009), values consistent with a chaotropic mode-of-action for a stressor with a $\log P$ of 1.36 (see Bhaganna *et al.*, 2010). *Saccharomyces cerevisiae* (like *P. anomala*, *Acinetobacter* and lactic acid bacteria) also inhibits the growth of competing species by acidifying the environment which it achieves via the production of a number of organic acids (Table 6; Kotyk *et al.*, 1999; Lowe and Arendt, 2004; Gulati *et al.*, 2010); it also secretes peptides with fungicidal and fungistatic activities (Albergaria *et al.*, 2010); and some strains contain inheritable virus-like particles containing genes that code for proteins inhibitory to other strains of the same species (Pretorius, 2000). Not only does *S. cerevisiae* employ multiple types of antimicrobial substance that act in concert, it is also has a correspondingly wide range of stressor-protection measures that minimize metabolic inhibition by antimicrobial stressors (those synthesized by either *S. cerevisiae* or other species), including efflux pumps (Pretorius, 2000),

and highly effective compatible-solute strategies (see above), protein-stabilization proteins and adaptations to acid stress (Table 6) and is therefore highly tolerant towards chaotropic and hydrophobic stressors (Fig. 2; Tables 4 and 6). *Saccharomyces cerevisiae* acid tolerance (as well as that of lactic acid bacteria) is conferred by an array of mechanisms (Table 3) that collectively contribute to the weed phenotype by removing the hydrogen ions and/or by promoting the refolding of partially denatured biomacromolecules.

Pichia anomala utilizes an array of antimicrobials that have synergistic effects as inhibitors and collectively target a wide array of taxa (including fungi, other yeasts, bacteria and viruses; Walker, 2011): chaotropic stressors such as ethanol and ethyl acetate; hydrophobic stressors such as isoamyl acetate; biosurfactants (see below); organic acids; an $\text{exo-}\beta\text{-1,3-glucanase}$ known to degrade glucans in fungal cell walls; and diverse toxins that have specific modes-of-action (see Table 6; Passoth *et al.*, 2006; Walker *et al.*, 2011). A semi-quantitative analysis of the antifungal activities of *P. anomala* suggests that inhibition results primarily from a combination of (in order of importance): killer toxins, hydrolytic enzymes and volatile substances (Table 6), as well as more minor factors such as nutrient competition and substrate acidification (Walker, 2011). *Rhodotorula mucilaginosa* produces an array of volatile chaotropic and hydrophobic stressors as well as killer toxins (Table 6). Many fungi also produce mycotoxins; for example *Aspergillus ochraceus* growing on maize grain can produce ochratoxins at up to 1.4 mg g^{-1} (Magan *et al.*, 2003). Indeed a number of *Aspergillus* species are notorious for the potent activity of their ochratoxins against other fungal strains and as well as bacteria (Table 6).

Like *S. cerevisiae*, some other microbial species are able to produce moderately chaotropic antimicrobials in bulk quantities: *Clostridium acetobutylicum* produces high concentrations of butanol, ethanol and acetone in anaerobic habitats, and *Zymomonas mobilis* produces high concentrations of ethanol (Tables S1 and 6). Butanol is a particularly potent chaotrope that is highly inhibitory at millimolar concentrations (Bhaganna *et al.*, 2010; Ezeji *et al.*, 2010; Cray *et al.*, 2013). We suspect that *Clostridium* spp. that can dominate chaotropic environments such as ammonia-rich animal slurries (Table S1) have an unusually high resistance to the net chaotropicity of the ammonia, ethanol, butanol, acetone and other chaotropic substances in their habitat (a hypothesis that has not yet been investigated to our knowledge; Williams and Hallsworth, 2009; Cray *et al.*, 2013). The chaotropicity of urea and ammonia is greater than that of ethanol when compared at equivalent molar concentrations (Hallsworth *et al.*, 2003a; Williams and Hallsworth, 2009) and there is evidence that some bacteria that secrete these compounds, including *Clostridium aminophilum* and

Pseudomonas stutzeri, benefit from chaotropicity-induced inhibition of their competitors (Table 6). There is no absolute resistance to the inhibitory/lethal activities of chaotropic and hydrophobic stressors (Sikkema *et al.*, 1995; Hallsworth, 1998; Hallsworth *et al.*, 2003a; 2003b; 2007; Duda *et al.*, 2004; Bhaganna *et al.*, 2010; Bell *et al.*, 2013) so the production of such antimicrobials coupled with reasonably effective self-protection mechanisms (see Tables S1 and 6) can sometimes be the ultimate means to achieve community dominance (see also Table S1).

Pseudomonas putida can generate and/or secrete diverse antimicrobial substances including catabolites produced from degradation of macromolecular substrates, VOCs and biosurfactants (Table 6). Biosurfactants that solubilize hydrocarbons and other hydrophobic substrates can also be highly inhibitory to other microbial species. Studies of *Pseudomonas* rhamnolipids suggest a key role in competitive ability (Wecke *et al.*, 2011); surfactants not only disorder plasma membranes and solubilize lipids, but can be chaotropic even for macromolecules that lack hydrophobic domains (Table 2; Cray *et al.*, 2013). The membrane-permeabilizing activities of rhamnolipids are particularly detrimental to Gram-positive bacteria, and are lethal to *Phytophthora* zoospores (Kruijt *et al.*, 2009; Wecke *et al.*, 2011). Bacteriocins produced by pseudomonads, like those of lactic acid bacteria, modify cellular membranes increasing permeability, causing leakage, disrupting transport processes and causing a dissipation of proton-motive force (Table 6). Catabolites such as phenol, catechol, vanillin, biphenyl, and sodium benzoate that are produced from the degradation of high-*M*_w hydrocarbons have inhibitory activity as chaotropic and hydrophobic stressors (Table 6; Hallsworth *et al.*, 2003a; Bhaganna *et al.*, 2010). *Pseudomonas putida* and other pseudomonads produce a wide variety of VOCs that collectively impact microbial competitors (Table 6). As outlined above, *P. putida* has multiple layers of defence against the potential stressful impact of solvent, chaotropic, and hydrophobic stressors via a versatile compatible solute metabolism (e.g. trehalose; Park *et al.*, 2007; Bhaganna *et al.*, 2010), protein-stabilization proteins, and production of extracellular polymeric substances etc. (Tables 3 and 6). In addition, and arguably the most important feature for stressor tolerance, *P. putida* cells are equipped with efflux pumps (for example the plasmid-encoded TtGHI) that are highly effective at expulsion of hydrocarbons such as toluene as well as many other stressors and toxins (Ramos *et al.*, 2002; Fillet *et al.*, 2012).

Whereas the aquatic habitats of planktonic species such as *D. salina*, *G. semen* and *M. aeruginosa* may serve to dilute secreted antimicrobials, there is evidence that VOCs produced by *D. salina*, *M. aeruginosa*, and other aquatic species (including compounds such as

β-cyclocitral and dichloromethane) are inhibitory to competing species (Table 6). *Gonyostomum semen* is able to lyse the cells of competing algal species on contact although the mechanism by which it does so has yet to be resolved (Table 6); and *M. aeruginosa* synthesizes a diverse range of volatile antimicrobials including phenazines and phenazine derivatives (Table 6; Lifshits and Carmeli, 2012). Lactic acid bacteria, in addition to bacteriocins, produce significant quantities of organic acids and secrete a vast array of inhibitory VOCs (Table 6). The analysis of volatiles produced, primarily by *L. rhamnosus*, during the production of a ewe's-milk cheese (Table S1) gives an insight into the complexity of the microbial warfare that takes place in open habitats: 57 compounds were detected including terpenes, alcohols, ketones and esters (Randazzo *et al.*, 2010). Although it is plausible that some of these substances exert specific toxic effects, all of them have the properties of either chaot- or hydrophobic stressors in microbial cells (Table 6; Bhaganna *et al.*, 2010), and their roles in habitat dominance has rarely received attention.

Microbial weed ecology

Open habitats

The concept of an open habitat is a fundamental, well-established principle in the fields of plant and animal ecology. For microorganisms, open habitats can be defined according to the diversity, interactions and community succession of their inhabitants in contrast to other types of microbial habitat that are defined by physicochemical characteristics, substrate type or nutrient concentration (Table 7). Hydrocarbons are the primary carbon substrate in approximately one-third of the open habitats listed in Table S1, and are generally derived from dense, non-aqueous phase liquids (e.g. oil) or plant metabolites secreted into the rhizosphere, phyllosphere or atmosphere (bioaerosols). A distinguishing feature of hydrocarbon-based systems is that they can remain perpetually open because, as dissolved hydrocarbons are assimilated, the substrate reservoir may be continually replenished by molecules from the oil entering the aqueous phase or (for the rhizosphere and phyllosphere) because the plant continues to produce and secrete hydrocarbons and other organics (Nunes *et al.*, 2005; Kazda and Falkinham III, 2009; McCammick *et al.*, 2009; Hunter *et al.*, 2010; Kachalkin and Yurkov, 2012). These habitats are commonly dominated by species such as *Acinetobacter johnsonii*, *Alkanivorax borkumensis*, *Pantoea ananatis* and *P. putida*. Whereas these organisms – all members of the gammaproteobacteria – may share many molecular and ecophysiological characteristics (see Tables S1 and 3–6) in most cases it is *P. putida* that emerges as the most prevalent species (Table S1).

Table 7. Properties of open (microbial) habitats.

	Value as a descriptor for an open habitat ^a	Notes and examples	
Typical characteristic(s)		For open habitats	For habitats that are not open
Microbial ecology			
Open to population by a multitude of microbial species	Primary	Potentially habitable by an indefinite number of species which may be from diverse Kingdoms of life (e.g. solar salterns see Table S1)	The fracture water of a South-African goldmine (2.8 km deep) was found to contain a single-species ecosystem (Chivian <i>et al.</i> , 2008)
Promote intense competition between microbial species	Primary	Conditions favour strong growth of many species and may select for ability to produce antimicrobial substances (see below)	A climax community, single-species habitat, and symbiotic partnership are characterized by low levels of inter-species competition
Can facilitate dominance of the developing community by one or more species (though this may not always take place)	Primary	Diverse types of substrate/environment can accommodate community progression towards dominance by one or more microbial species (see Table S1)	The antithesis of an open habitat is one that is occupied by a climax community that is no longer capable of further evolution (e.g. Paul <i>et al.</i> , 2006)
Transient and self-terminating; unless maintained in an 'open' condition	Primary	Frequently leading to the development of a closed community dominated by a single species (see Fig. 1; Table S1); unless 're-opened' due to grazing for example, or replenished (e.g. by fresh inputs of nutrients)	Closed habitats typically lack conditions that facilitate biomass formation and species succession
Resident microbes			
Strong selection for microbes able to out-grow competitors	Primary	For example, habitats rich in nutrients may favour species of bacterial copiotroph which may merge as dominating species if they possess other weed traits	A habitat with limited available resources would not would not support growth (the latter is associated with competition/ species succession)
Open to manipulation by species able to modify environmental conditions and/or secrete antimicrobials	Primary	Dynamic biomass formation makes open habitats available for/ vulnerable to manipulation by species that can introduce growth-limiting substances (Table 6; Cordero <i>et al.</i> , 2012)	In closed habitats the community is already restricted in terms of space, nutrients and/or physicochemical constraints etc.
Select for microbes that lack specific requirements (such as rare trace elements, or a specific host)	Secondary	Microbes such as <i>S. cerevisiae</i> and <i>Lactobacillus</i> spp. typically grow without idiosyncratic requirements (such as a requirement for interactions with other species)	Specialist microbes are found in niche habitats, such as those that form lichens (Hoffland <i>et al.</i> , 2004)
Chemistry and properties of the cellular environment			
Not excessively stressful for resistant microbes in terms of water activity, chaotropicity, pH, temperature etc. ^b	Primary	Open (microbial) habitats are, by definition, open to population by a large number and high diversity of microbial species so are typically not physically or chemically extreme ^c	Extreme habitats that are occupied by a small number of obligate extremophiles rarely support the high growth rates or promote the intense competition that typify open habitats ^c
Have the potential to support a substantial microbial biomass and so do not constrain ecosystem development	Primary	Open habitats have sufficient space and resources to facilitate the ecosystem development (Table S1; Fig. 1)	Non-open habitats can be constrained by nutritional and/or physicochemical parameters (Chivian <i>et al.</i> , 2008)
Not contaminated with inhibitory substances such as heavy metals or other potent toxins	Secondary	Some polluted, hydrocarbon-rich environments may also support the development of microbial communities	High levels of toxic substances will prevent high growth rates and minimize diversity

a. Those designated as descriptors of primary importance are considered essential to the definition of an open (microbial) habitat.

b. Not so extreme that there is limited diversity or possibility for biomass formation: i.e. typically > 0.75 water activity (see Pitt, 1975; Williams and Hallsworth, 2009), < 20–25 kJ kg⁻¹ mole⁻¹ chaotropic activity (Hallsworth *et al.*, 2007), < pH 11 or > -15°C.

c. Whereas NaCl-saturated habitats are generally considered extreme, their water activity is ³ 0.75 – well above that which ultimately limits microbial life (i.e. 0.61–0.71: Pitt and Christian, 1968; Brown, 1990; Grant, 2004; Williams and Hallsworth, 2009; Leong *et al.*, 2011) – and typically supports a remarkably high microbial diversity and biomass (Antón *et al.*, 2002; Daffonchio *et al.*, 2006; Baati *et al.*, 2008; Bardavid *et al.*, 2008; Khemakhem *et al.*, 2010).

This is a product of its metabolic versatility, exceptional tolerance to hydrocarbons and other solutes, and antimicrobial activities (Tables 3, 5 and 6). Other weed species, however, have greater acid tolerance than the gammaproteobacteria: in the acidic surface film of the sphagnum-moss grey layer it is *Mycobacterium* (including *Mycobacterium terrae*, *M. smegmatis* and *Mycobacterium sphagnum*) that emerges as a dominant genus; and in the green layer at lower pH (2–2.5), *R. mucilaginosa* has the competitive advantage (Table S1). Phyllosphere, rhizosphere and other habitats can also remain continuously open if they are replenished with organic substances from ions released from soil particles, extracellular polymeric substances and compatible solutes from microbial cells, root diffusates, animal wastes, diverse sources of necromass, etc.

Habitats which facilitate weed emergence are typically not so extreme that there is a negligible microbial diversity or little capacity for community development/biomass formation; i.e. > 0.75 water activity, < 20–25 kJ kg⁻¹ mole⁻¹ chaotropic activity, < pH 11 or > –15°C (Table 7). Open habitats have available nutrients, electron acceptors etc and C : N ratios that favour microbial growth, and do not contain high levels of inhibitory substances such as heavy metals or toxic pollutants. They will therefore promote intense competition between species assuming that a diversity of microbial species is present (Tables S1 and 7). Whereas high-sugar habitats with a water activity below 0.9 are too stressful for the vast majority of microbes, including *S. cerevisiae*, there is still a considerable diversity of xerotolerant and xerophilic species that are highly active between water activities of 0.84 and 0.9 (Pitt, 1975; Brown, 1990). Similarly, crystallizer ponds or solar salt-erns are known as extreme environments and only a small fraction of microbial species can survive and grow at saturated NaCl. Nevertheless, it is the limited solubility of NaCl (and not the limitations of salt-adapted macromolecular and cellular systems) that determines the limit for microbial growth in high-NaCl habitats: a number of obligate halophiles grow optimally at 35% w/v NaCl (0.755 water activity) and different types of NaCl-saturated environment are biodiverse, highly competitive, biomass-rich microbial habitats (Antón *et al.*, 2002; Daffonchio *et al.*, 2006; Baati *et al.*, 2008; Bardavid *et al.*, 2008; Khemakhem *et al.*, 2010). Salterns (and also algal mats) may under some conditions function as continually open habitats if nutrients are not limiting; *D. salina* for example can release sufficient glycerol into the system to satisfy the carbohydrate requirement of the wider community (Borowitzka, 1981; Bardavid *et al.*, 2008).

Open habitats favour microbes with shorter generation-times and relatively vigorous growth-phenotypes and in this way may drive the evolution of faster-growing and more stress-tolerant species (as well as those with effec-

tive antimicrobial strategies; Tables 4, 6 and 7). That the ecology and evolution of plant weeds are intimately and inextricably linked with open habitats has been established for some time (Anderson, 1952); in contrast there is little information about open habitats of microorganisms (see Table 1; Hallsworth, 1998). Open habitats can be spontaneously created (e.g. due to removal of microbial biomass by predators and grazers; McArthur, 2006), dynamic (e.g. due to desiccation–rehydration cycles or nutrient limitation), and/or transient (Tables S1 and 7). As is true for open habitats of plants (Anderson, 1952), open habitats of microbes can also become closed if they become water-constrained or otherwise resource-depleted; and/or the microbial community reaches stationary phase (Paul *et al.*, 2006): and these changes may be accompanied by – or even a consequence of – domination by weed species. Environments can even progress to become sterile; for example those with high ethanol concentrations (Hallsworth, 1998; Pretorius, 2000). However, a dualistic concept of open habitats on the one hand and closed habitats on the other would be simplistic; in reality there is a seamless continuum between the two extremes of fertile but uncolonized and hostile/uninhabitable environment. Habitats with a favourable water activity for most bacteria (≥ 0.94) and high concentrations of carbon and nitrogen substrates such as milk, cheese and crushed peanuts (Table S1) typically progress to a closed condition (and are dominated by lactic acid bacteria), as do most high-sugar habitats (Table S1). A number of the open habitats listed in Table S1 that are resource-rich, and stressful, temporally or spatially heterogenous, and/or require saprotrophic nutrition were dominated by *Pichia* (e.g. moist barley grains; salted fermenting olives) or *Aspergillus* species (e.g. decomposing leaf litter, sugarcane juice). The stress biology, energy-generating capabilities and antimicrobial activities of these species undoubtedly contribute to their success (Tables 3 and 6).

Ecophysiological classification

Microorganisms have been previously categorized based on the way in which their phenotype impacts ecological behaviour: as autotrophs and heterotrophs; generalists and specialists; *r*- and *K*-strategists, aerobes and anaerobes; copiotrophs and oligotrophs; fast and slow growers; and extremophiles and mesophiles (see Table 4 and S2). The definition of microbial weeds is qualitatively distinct from that of other microbial groups (see Table S2): only weeds are characterized by their ability to dominate communities and the associated/underlying biology (Tables S1, 2–6 and S2) and it is only weeds that are linked via their ecology and evolution with open habitats (Tables 1, S1, 3 and S2). The biology and ecology of microbial weed species cuts across established groupings of

microorganisms (i.e. the former does not represent a parallel category; see Table S2). Some microbes have a considerable number of weed traits but only rarely emerge as dominant members of communities (e.g. *Pseudoxylaria* spp., *E. coli* and *M. smegmatis*; see Tables S1 and 3, 5 and 6). Such species serve as a reminder that weeds and non-weeds represent two extremes of an otherwise continuous scale. An insightful study of *Termitomyces* ecology, for example, discusses the biology of *Pseudoxylaria*; a fungus that is present in *Termitomyces* habitats but is typically inactive and/or present at negligible cell densities (Visser *et al.*, 2011). *Pseudoxylaria* could be ecologically disadvantaged if present in greater numbers as termites, which can detect VOCs produced during interactions between *Pseudoxylaria* and *Termitomyces*, remove the unwanted fungus from their *Termitomyces* cultures. The extant *Pseudoxylaria* strains thereby appear to have been selected for/have evolved a low competitive ability (Visser *et al.*, 2011). Nevertheless *Pseudoxylaria* can occasionally emerge to dominate the substrate (Visser *et al.*, 2011), a fertile open habitat consisting of decomposing plant material (Table S1). Non-weed species make up a loose affiliation of species that are only linked by their poor ability to dominate communities in open habitats and may not share any specific metabolic or phenotypic traits (Table S2). We propose that this 'dustbin' group includes *Synodropsis* spp., *Polypaecilum pisce*, *Metschnikowia orientalis*, *Caulobacter crescentus* and *Salmonella* species (Table S2). Several ecophysiological classification systems that are based on microbial growth-phenotype and/or substrate utilization (Table S2). Microbial generalists can occupy a wide range of ecological niches, and typically achieve this via the ability to utilize a wide range of nutrients and substrates (Table S2). Conversely microbial specialists, such as *S. cerevisiae*, *H. werneckii* and many basidiomycetes have specific nutritional and/or physicochemical requirements (de Fine Licht *et al.*, 2005; Kashangura *et al.*, 2006). Obligate extremophiles commonly occupy, and are adapted to grow optimally in, niches that are hostile to the majority of microorganisms (Table S2). There have been relatively few studies (and is consequently inadequate terminology) to properly describe microbes that are extremophile-intolerant such as xero-intolerant species (see Table 4), with the exception of mesophilic species that function optimally at non-extreme temperatures (though the term *mesophiles* can be used in a general sense to refer to non-extremophiles: Table S2; Gostinčar *et al.*, 2010). Non-extremophilic microbes grow optimally in physicochemical conditions which favour metabolic activity and growth of the majority of microbial species, and this grouping includes both weed and non-weed species (Table S2). While fast-growing microbes have short doubling times, the vast majority of these species do not

have weed-like capabilities and are unable to out-grow their competitors (Tables 4 and S2). It may be that relatively few oligotrophic microbes or *K*-strategists are weed species, although some mycobacteria do have a weed phenotype (Tables S1, 3, 5 and 6). *Pseudomonas putida* is generally considered to be a model *r*-strategist (e.g. Margesin *et al.*, 2003) yet *P. putida* has an extraordinary competitive ability (see Tables S1 and 4–6), and may not be the best example of an *r*-strategist (the latter are characterized by a weak competitive ability; Table S2). By contrast *P. putida* appears to be an exceptional example of a microbial weed both in terms of its cellular biology and its ability to dominate diverse types of open habitat.

Concluding remarks

It is irrefutable that a small minority of microbial species recurrently appear as the dominant members of microbial communities; and we suggest here that these microbes have shared phenotypic traits. This implies that open-habitat ecology has impacted the evolutionary trajectories of these microbes, and that the activities of weeds determine microbial diversity in many localities within the biosphere. Whereas microbial weed species are not always useful to humankind (see Table 1), they make up a substantial portion of the world's biotechnology sector. *Saccharomyces cerevisiae* supports global industries in fermented products, biofuel, biocides (ethanol is also an important disinfectant; McDonnell and Russell, 1999), and other white biotechnologies; *P. anomala* produces a substantial catalogue of biotechnologically useful substances and is employed in production of foodstuffs (see Walker, 2011); lactic acid bacteria form the basis of food- and drinks-production processes where the antimicrobial volatiles and organic acids that they secrete act as flavour compounds and natural preservatives, and are used as probiotics, and for the production of fine chemicals (Rose, 1982; Steinkraus, 1983; Molly *et al.*, 1996; Nomura *et al.*, 1998; Tiwari *et al.*, 2009; Randazzo *et al.*, 2010; Bron *et al.*, 2012); *P. putida* is used for the commercial production of bulk and fine chemicals, industrial biocatalysis, and as the basis of many other industrial and environmental biotechnologies (Kruijt *et al.*, 2009; Puchalka *et al.*, 2008; Hassan *et al.*, 2011; Poblete-Castro *et al.*, 2012); a number of weed species, including *Aspergillus* spp., *P. putida* and *Clostridium* spp. play primary roles in the treatment or decomposition of plant matter, oil and hydrocarbons, xenobiotics in polluted soils, sludge and slurry treatments, and other organic wastes (Tables S2 and 4; Narihiro and Sekiguchi, 2007; Timmis, 2009); and *Aspergillus* species are used for production of foods (Rose, 1982; Tamang, 2010), fine chemicals (e.g. citric acid, secondary metabolites) and extracellular enzymes

(e.g. amylase, cellulase, pectinase and protease), for industrial biocatalysis, and have potential for bioremediation and biosorption applications (Ward *et al.*, 2006). It is the metabolic activities of substances that suppress potential competitors which make *S. cerevisiae*, *Rhodotorula* spp., *P. anomala*, *P. putida*, lactic acid bacteria and other weed species invaluable as agents of biological control, for preventing spoilage of drinks, foodstuffs and animal feeds (see Calvente *et al.*, 1999; Kruijt *et al.*, 2009; Hassan *et al.*, 2011; Olstorpe and Passoth, 2011; Walker, 2011).

Microbial weeds are also used as model systems for research, yet key aspects of their biology remain enigmatic. Many plant weeds have exceptional reproduction and dispersal systems (Table 1) and it remains unclear whether microbial weed species produce more propagules, have superior dispersal mechanisms, or evolved propagules with exceptional longevity, and/or the ability to germinate over a wider range of conditions than their non-weed comparators. Could it be that, like plant weeds, the competitive ability of microbial weeds is enhanced by an abnormally high phenotypic plasticity and intra-specific variability (see Table 1)? Out of the world's plant flora it has been estimated that only several hundred plant species are weeds (Hill, 1977); what percentage of microbes might exhibit weed phenotypes and behaviour? Is there evidence preserved in rocks, sediments, peat, ice, or the hypersaline fluid inclusions of salt crystals that can shed light on the evolutionary trajectories of microbial weed species? Is there an ideal genome size for prokaryotic and eukaryotic weed species? What are the minimum/essential characteristics required to enable a microorganism to dominate a specific habitat; can we design and manufacture a microbial weed via a synthetic biology approach? Will hitherto uninvestigated microbial weeds prove to be rich sources of antimicrobial compounds for novel biotechnological and clinical applications? Can microbial weed ecology shed light on wound microbiology, lead to the prevention of the microbial colonization of prosthetic devices, or minimize the emergence of opportunistic pathogens? Could knowledge-based interventions in weed ecology enhance our ability to manipulate and manage microbes at various levels of food-production and food-spoilage, or lead to better management of environments that harbor potentially dangerous species?

Crop plants such as sunflowers, potatoes, wheat and barley are known to have weedy ancestors (Anderson, 1952) and it is thought that the evolution of both plant weeds and crop plants has been driven by anthropogenically opened habitats. We believe that open (microbial) habitats also select for the phenotypes, and therefore act as a potent driving force that determines the evolutionary trajectories, of microbial weeds and their communities:

open-habitat ecology may therefore be worthy of recognition as a distinct scientific field that spans all organismal biology including microbiology. There is a maxim that everything is everywhere but that the environment selects; this implies that microbial species are everywhere on Earth but that the local environment determines which are able to grow (Baas Becking, 1934; de Wit and Bouvier, 2006). While not necessarily untrue, this idea implies a level of passivity on the part of the microbial cell. Yet microorganisms are inherently variable and dynamic systems with considerable genetic dexterity and phenotypic plasticity (able to both respond and adapt). Baas Becking (1934) acknowledged that cells are able to modify their extracellular environment, but microbes are also proactive in their manipulation of each other. It may be that microbial weeds are the most accomplished examples of this.

Acknowledgements

We are most grateful to T.A. Hill (University of London, UK) for his inspirational lectures on the biology of plant weeds; and for scientific enjoyable and fruitful discussions with E. J. Carpenter (San Francisco State University, USA), L.R. Cooke (Agri-Food and Biosciences Institute, Belfast); B.T.M. Dentinger (Royal Botanic Gardens, Kew, UK), N. Gunde-Cimerman (University of Ljubljana, Slovenia), S. Iwaguchi (Nara Women's University, Japan), M.-A. Lachance (University of Western Ontario, Canada), K.D. Farnsworth, I.R. Grant, J.D.R. Houghton, C.J. Law, A.L. McCann, J.W. McGrath, J. Megaw, J. P. Quinn, and J.T. Russell (Queen's University Belfast), J.L. Green, and A.M. Womack (University of Oregon, USA), T.J. McGenity (University of Essex, UK), T. Mock (University of East Anglia), A. Oren (The Hebrew University of Jerusalem, Israel), I.S. Pretorius (University of South Australia, Australia), R. Santos (Laboratório de Análises of the Instituto Superior Técnico, Lisbon, Portugal), R.S. Singhal (Institute of Chemical Technology, Mumbai, India), G.G. Stewart (Heriot-Watt University, UK), J.P. Tamang (Sikkim University, India), K.N. Timmis (Helmholtz Centre for Infection Research, Braunschweig, Germany), M.A. Voytek (NASA Headquarters, USA).

Conflict of interest

None declared.

References

- Albergaria, H., Francisco, D., Gori, K., Arneborg, N., and Gírio, F. (2010) *Saccharomyces cerevisiae* CCMI 885 secretes peptides that inhibit the growth of some non-*Saccharomyces* wine-related strains. *Appl Microbiol Biotechnol* **86**: 965–972.

- Albertyn, J., Hohmann, S., Thevelein, J.M., and Prior, B. (1994) *GPD1*, which encodes glycerol-3-phosphate dehydrogenase, is essential for growth under osmotic stress in *Saccharomyces cerevisiae*, and its expression is regulated by the high osmolarity glycerol response pathway. *Mol Cell Biol* **14**: 4135–4144.
- Alsaadawi, I.S., and Salih, N.M.M. (2009) Allelopathic potential of *Cyperus rotundus* L. I. Interference with crops. *Allelopathy J* **23**: 297–303.
- Al-Tahhan, R.A., Sandrin, T.R., Bodour, A.A., and Maier, R.M. (2000) Rhamnolipid-induced removal of lipopolysaccharide from *Pseudomonas aeruginosa*: effect on cell surface properties and interaction with hydrophobic substrates. *Appl Environ Microbiol* **66**: 3263–3268.
- Altermann, E., Russell, W.M., Azcarate-Peril, M.A., Barrangou, R., Buck, B.L., McAuliffe, O., et al. (2005) Complete genome sequence of the probiotic lactic acid bacterium *Lactobacillus acidophilus* NCFM. *Proc Natl Acad Sci USA* **102**: 3906–3912.
- Álvarez-Ordóñez, A., Begley, M., Prieto, M., Messens, W., López, M., Bernardo, A., et al. (2011) *Salmonella* spp. survival strategies within the host gastrointestinal tract. *Microbiology* **157**: 3268–3281.
- Andersen, D., Renshaw, J.C., and Wiebe, M.G. (2003) Rhodotorulic acid production by *Rhodotorula mucilaginosa*. *Mycol Res* **107**: 949–956.
- Anderson, E. (1952) *Plants, Man and Life*. Boston, MA, USA: Little, Brown and Company.
- Anderson, R.C., Flythe, M.D., Krueger, N.A., Callaway, T.R., Edrington, T.S., Harvey, R.B., et al. (2010) Decreased competitiveness of the foodborne pathogen *Campylobacter jejuni* during co-culture with the hyper-ammonia producing anaerobe *Clostridium aminophilum*. *Folia Microbiol* **55**: 309–311.
- Anderssen, E.L., Diep, D.B., Nes, I.F., Eijsink, V.G.H., and Nissen-Meyer, J. (1998) Antagonistic activity of *Lactobacillus plantarum* C11: two new two-peptide bacteriocins, plantaricins EF and JK, and the induction factor plantaricin A. *Appl Environ Microbiol* **64**: 2269–2272.
- Antón, J., Oren, A., Benlloch, S., Rodríguez-Valera, F., Amann, R., and Rosselló-Mora, R. (2002) *Salinibacter ruber* gen. nov., sp. nov., a novel, extremely halophilic member of the *Bacteria* from saltern crystallizer ponds. *Int J Syst Evol Microbiol* **52**: 485–491.
- Aranda, J.S., Salgado, E., and Taillandier, P. (2004) Trehalose accumulation in *Saccharomyces cerevisiae* cells: experimental data and structured modeling. *Biochem Eng J* **17**: 129–140.
- Arsène, F., Tomoyasua, T., and Bukau, B. (2000) The heat shock response of *Escherichia coli*. *Int J Food Microbiol* **55**: 3–9.
- Aziz, S., Memon, H.R., Shah, F.A., Rajoka, M.I., and Soomro, S.A. (2009) Production of ethanol by indigenous wild and mutant strain of thermotolerant *Kluyveromyces marxianus* under optimized fermentation conditions. *Pak J Anal Environ Chem* **10**: 25–33.
- Baas Becking, L.G.M. (1934) *Geobiologie of inleiding tot de milieukunde*. The Hague, the Netherlands: W.P. Van Stockum 0026; Zoon (in Dutch).
- Baati, H., Guermazi, S., Amdouni, R., Gharsallah, N., Sghir, A., and Ammar, E. (2008) Prokaryotic diversity of a Tunisian multipond solar saltern. *Extremophiles* **12**: 505–518.
- Back, J.F., Oakenfull, D., and Smith, M.B. (1979) Increased thermal-stability of proteins in the presence of sugars and polyols. *Biochemistry* **18**: 5191–5196.
- Baker, H.G. (1965) Characteristics and modes of origin of weeds. In *Genetics of Colonizing Species*. Baker, H.G., and Stebbins, G.L. (eds). New York, USA: Academic Press, pp. 147–168.
- Bakker, E.G., Montgomery, B., Nguyen, T., Eide, K., Chang, J., Mockler, T.C., et al. (2009) Strong population structure characterizes weediness gene evolution in the invasive grass species *Brachypodium distachyon*. *Mol Ecol* **18**: 2588–2601.
- Balashov, S.P., Imasheva, E.S., Boichenko, V.A., Anton, J., Wang, J.M., and Lanyi, J.K. (2005) Xanthorhodopsin: a proton pump with a light-harvesting carotenoid antenna. *Science* **309**: 2061–2064.
- Ballerstedt, H., Volkers, R.J.M., Mars, A.E., Hallsworth, J.E., Martins dos Santos, V.A., Puchalka, J., et al. (2007) Genotyping of *Pseudomonas putida* strains using *P. putida* KT2440-based high-density DNA microarrays: implications for transcriptomics studies. *Appl Microbiol Biotechnol* **75**: 1133–1142.
- Bardavid, R.E., and Oren, A. (2008) Sensitivity of *Haloquadratum* and *Salinibacter* to antibiotics and other inhibitors: implications for the assessment of the contribution of Archaea and Bacteria to heterotrophic activities in hypersaline environments. *FEMS Microbiol Ecol* **63**: 309–315.
- Bardavid, R.E., and Oren, A. (2012) The amino acid composition of proteins from anaerobic halophilic bacteria of the order Halanaerobiales. *Extremophiles* **16**: 567–572.
- Bardavid, R.E., Khristo, P., and Oren, A. (2008) Interrelationships between *Dunaliella* and halophilic prokaryotes in saltern crystallizer ponds. *Extremophiles* **12**: 5–14.
- Behrends, V., Williams, K.J., Jenkins, V.A., Robertson, B.D., and Bundy, J.G. (2012) Free glucosylglycerate is a novel marker of nitrogen stress in *Mycobacterium smegmatis*. *J Proteome Res* **11**: 3888–3896.
- Bell, A.N.W., Magill, E., Hallsworth, J.E., and Timson, D.J. (2013) Effects of alcohols and compatible solutes on the activity of β -galactosidase. *Appl Biochem Biotechnol*. doi: 10.1007/s12010-012-0003-3.
- Bellinger, B.J., and Van Mooy, B.A.S. (2012) Nonphosphorus lipids in *Periphyton* reflect available nutrients in the Florida Everglades, USA. *J Phycol* **48**: 303–311.
- Benning, C. (1998) Biosynthesis and function of the sulfolipid sulfoquinovosyl diacylglycerol. *Annu Rev Plant Phys* **49**: 53–75.
- Bental, M., Pick, U., Avron, M., and Degani, H. (1991) Polyphosphate metabolism in the alga *Dunaliella salina* studied by P-31-NMR. *Biochim Biophys Acta* **1092**: 21–28.
- Berdjeb, L., Pollet, T., Domaizon, I., and Jacquet, S. (2011) Effect of grazers and viruses on bacterial community structure and production in two contrasting trophic lakes. *BMC Microbiol* **11**: Article number 88.
- Bernal, P., Seguar, A., and Ramos, J.-L. (2007) Compensatory role of the *cis-trans*-isomerase and cardiolipin

- synthase in the membrane fluidity of *Pseudomonas putida* DOT-T1E. *Environ Microbiol* **9**: 1658–1664.
- Berrah, G., and Konetzka, W.A. (1962) Selective and reversible inhibition of the synthesis of bacterial deoxyribonucleic acid by phenethyl alcohol. *J Bacteriol* **83**: 738–744.
- Berry, C., Fernando, W.G.D., Loewen, P.C., and de Kievit, T.R. (2010) Lipopeptides are essential for *Pseudomonas* sp. DF41 biocontrol of *Sclerotinia sclerotiorum*. *Biol Control* **55**: 211–218.
- Bhaganna, P., Volkers, R.J.M., Bell, A.N.W., Kluge, K., Timson, D.J., McGrath, J.W., et al. (2010) Hydrophobic substances induce water stress in microbial cells. *Microb Biotechnol* **3**: 701–716.
- Bolhuis, H., Palm, P., Wende, A., Falb, M., Rampp, M., Rodriguez-Valera, F., et al. (2006) The genome of the square archaeon *Haloquadratum walsbyi*: life at the limits of water activity. *BMC Genomics* **7**: 169.
- Bonnet, M.P., and Poulin, M. (2002) Numerical modelling of the planktonic succession in a nutrient-rich reservoir: environmental and physiological factors leading to *Microcystis aeruginosa* dominance. *Ecol Modell* **156**: 93–112.
- Borowitzka, L.J. (1981) The microflora. Adaptation to life in extremely saline lakes. *Hydrobiologia* **81**: 33–46.
- Boutibonnes, P., Auffray, Y., Malherbe, C., Kogbo, W., and Marais, C. (1984) Antibacterial and genotoxic properties of 33 mycotoxins. *Mycopathologia* **87**: 43–49.
- Boynton, Z.L., Bennett, G.N., and Rudolph, F.B. (1996) Cloning, sequencing, and expression of genes encoding phosphotransacetylase and acetate kinase from *Clostridium acetobutylicum* ATCC 824. *Appl Environ Microbiol* **62**: 2758–2766.
- Britigan, B.E., Miller, R.A., Hassett, D.J., Pfaller, M.A., McCormick, M.L., and Rasmussen, G.T. (2001) Antioxidant enzyme expression in clinical isolates of *Pseudomonas aeruginosa*: Identification of an atypical form of manganese superoxide dismutase. *Infect Immun* **69**: 7396–7401.
- Bron, P.A., Van Baaren, P., and Kleerebezem, M. (2012) Emerging molecular insights into the interaction between probiotics and the host intestinal mucosa. *Nat Rev Microbiol* **10**: 66–U90.
- Brown, A.D. (1990) *Microbial Water Stress Physiology – Principles and Perspectives*. Chichester, UK: Wiley.
- Buzzini, P., Gasparetti, C., Turchetti, B., Cramarossa, M.R., Vaughan-Martini, A., Martini, A., et al. (2005) Production of volatile organic compounds (VOCs) by yeasts isolated from the ascocarps of black (*Tuber melanosporum* Vitt.) and white (*Tuber magnatum* Pico) truffles. *Arch Microbiol* **184**: 187–193.
- Cabrol, L., Malhautier, L., Poly, F., Lepeuple, A.-S., and Fanlo, J.-L. (2012) Bacterial dynamics in steady-state biofilters: beyond functional stability. *FEMS Microbiol Ecol* **79**: 260–271.
- Calvente, V., Benuzzi, D., and de Tossati, M.I.S. (1999) Antagonistic action of siderophores from *Rhodotorula glutinis* upon the postharvest pathogen *Penicillium expansum*. *Int Biodeterior Biodegradation* **43**: 167–172.
- Capone, K.A., Dowd, S.E., Stamatas, G.N., and Nikolovski, J. (2011) Diversity of the human skin microbiome early in life. *J Invest Dermatol* **131**: 2026–2032.
- Casadei, M.A., Mañas, P., Niven, G., Needs, E., and Mackey, B.M. (2002) Role of membrane fluidity in pressure resistance of *Escherichia coli* NCTC 8164. *Appl Environ Microbiol* **68**: 5965–5972.
- Castrol, C.D., Koretsky, A.P., and Domach, M.M. (1999) NMR-observed phosphate trafficking and polyphosphate dynamics in wild-type and vph1-1 mutant *Saccharomyces cerevisiae* in response to stresses. *Biotechnol Prog* **15**: 65–73.
- Cerdeño-Tárraga, A.M., Patrick, S., Crossman, L.C., Blakely, G., Abratt, V., Lennard, N., et al. (2005) Extensive DNA inversions in the *B. fragilis* genome control variable gene expression. *Science* **307**: 1463–1465.
- Chakrabarty, A.M., and Roy, S.C. (1964) Nature of endogenous reserve material in a strain of *Pseudomonas fluorescens-putida* intermediate. *Biochem J* **92**: 105–112.
- Chang, W.-S., van de Mortel, M., Nielsen, L., de Guzman, G.N., Li, X., and Halverson, L.J. (2007) Alginate production by *Pseudomonas putida* creates a hydrated microenvironment and contributes to biofilm architecture and stress tolerance under water-limiting conditions. *J Bacteriol* **189**: 8290–8299.
- Chen, G., Bradford, W.D., Seidel, C.W., and Li, R. (2012) Hsp90 stress potentiates rapid cellular adaptation through induction of aneuploidy. *Nature* **482**: 246–250.
- Chen, H., and Jiang, J.-G. (2009) Osmotic responses of *Dunaliella* to the changes of salinity. *J Cell Physiol* **219**: 251–258.
- Chen, J., Lee, S.M., and Mao, Y. (2004) Protective effect of exopolysaccharide colanic acid of *Escherichia coli* O157:H7 to osmotic and oxidative stress. *Int J Food Microbiol* **93**: 281–286.
- Chen, M., Tang, H.Y., Ma, H.Z., Holland, T.C., Ng, K.Y.S., and Salley, S.O. (2011) Effect of nutrients on growth and lipid accumulation in the green algae *Dunaliella tertiolecta*. *Bioresour Technol* **102**: 1649–1655.
- Chin, J.P., Megaw, J., Magill, C.L., Nowotarski, K., Williams, J.P., Bhaganna, P., et al. (2010) Solutes determine the temperature windows for microbial survival and growth. *Proc Natl Acad Sci USA* **107**: 7835–7840.
- Chirife, J., Favetto, G., and Fontán, C.F. (1984) Microbial growth at reduced water activities: some physicochemical properties of compatible solutes. *J Appl Bacteriol* **56**: 259–268.
- Chiruvolu, V., Eskridge, K., Cregg, J., and Meagher, M. (1998) Effects of glycerol concentration and pH on growth of recombinant *Pichia pastoris* yeast. *Appl Biochem Biotechnol* **75**: 163–173.
- Chivian, D., Brodie, E.L., Alm, E.J., Culley, D.E., Dehal, P.S., DeSantis, T.Z., et al. (2008) Environmental genomics reveals a single-species ecosystem deep within Earth. *Science* **322**: 275–278.
- Chou, C.H. (2010) Role of allelopathy in sustainable agriculture: use of allelochemicals as naturally occurring bioagrochemicals. *Allelopathy J* **25**: 3–16.
- Coleman, A.W., and Heywood, P. (1981) Structure of the chloroplast and its DNA in chloromonadophycean algae. *J Cell Sci* **49**: 401–409.
- Collado-Fabbi, S., Vaulot, D., and Ulloa, O. (2011) Structure and seasonal dynamics of the eukaryotic picophytoplankton community in a wind-driven coastal upwelling ecosystem. *Limnol Oceanogr* **56**: 2334–2346.

- Colomb, A., Yassaa, N., Williams, J., Peeken, I., and Lochte, K. (2008) Screening volatile organic compounds (VOCs) emissions from five marine phytoplankton species by head space gas chromatography/mass spectrometry (HS-GC/MS). *J Environ Monit* **10**: 325–330.
- Corcelli, A., Lattanzio, V.M.T., Mascolo, G., Babudri, F., Oren, A., and Kates, M. (2004) Novel sulfonolipid in the extremely halophilic bacterium *Salinibacter ruber*. *Appl Environ Microbiol* **70**: 6678–6685.
- Corcuff, R., Mercier, J., Tweddell, R., and Arul, J. (2011) Effect of water activity on the production of volatile organic compounds by *Muscodor albus* and their effect on three pathogens in stored potato. *Fungal Biol* **115**: 220–227.
- Cordero, O.X., Wildschutte, H., Kirkup, B., Proehl, S., Ngo, L., Hussain, F., et al. (2012) Ecological populations of bacteria act as socially cohesive units of antibiotic production and resistance. *Science* **337**: 1228–1231.
- Costa, V., Amorim, M.A., Reis, E., Quintanilha, A., and Moradas-Ferreira, P. (1997) Mitochondrial superoxide dismutase is essential for ethanol tolerance of *Saccharomyces cerevisiae* in the post-diauxic phase. *Microbiology* **43**: 1649–1656.
- Couturier, E., and Rocha, E.P.C. (2006) Replication-associated gene dosage effects shape the genomes of fast-growing bacteria but only for transcription and translation genes. *Mol Microbiol* **59**: 1506–1518.
- Cox, R.A. (2004) Quantitative relationships for specific growth rates and macromolecular compositions of *Mycobacterium tuberculosis*, *Streptomyces coelicolor* A3(2) and *Escherichia coli* B/r: an integrative theoretical approach. *Microbiology* **150**: 1413–1426.
- Cramer, H.H. (1967) *Plant Protection and World Crop Production*. Pflanzenschutz: Nachrichten Bayer.
- Cray, J.A., Russell, J.T., Timson, D.J., Singhal, R.S., and Hallsworth, J.E. (2013) A universal measure of chaotropicity and kosmotropicity. *Environ Microbiol* **15**: 287–296.
- Crowe, J.H., Crowe, L.M., and Chapman, D. (1984) Preservation of membranes in anhydrobiotic organisms – the role of trehalose. *Science* **223**: 701–703.
- D'Amore, T., Panchal, C.J., Russell, I., and Stewart, G.G. (1990) A study of ethanol tolerance in yeast. *Crit Rev Biotechnol* **9**: 287–304.
- D'Souza-Ault, M.R., Smith, L.T., and Smith, G.M. (1993) Roles of N-acetylglutaminylglutamine amide and glycine betaine in adaptation of *Pseudomonas aeruginosa* to osmotic stress. *Appl Environ Microbiol* **59**: 473–478.
- Daffonchio, D., Borin, S., Brusa, T., Brusetti, L., van der Wielen, P.W.J.J., Bolhuis, H., et al. (2006) Stratified prokaryote network in the oxic–anoxic transition of a deep sea halocline. *Nature* **440**: 203–207.
- Daniels, C., Godoy, P., Duque, E., Molina-Henares, M.A., de la Torre, J., del Arco, J.M., et al. (2010) Global regulation of food supply by *Pseudomonas putida* DOT-T1E. *J Bacteriol* **192**: 2169–2181.
- De Fine Licht, H.H., Andersen, A., and Aanen, D.K. (2005) *Termitomyces* sp. associated with the termite *Macrotermes natalensis* has a heterothallic mating system and multinucleate cells. *Mycol Res* **109**: 314–318.
- Desai, J.D., and Banat, I.M. (1997) Microbial production of surfactants and their commercial potential. *Microbiol Mol Biol Rev* **61**: 47–64.
- Dhar, R., Sägesser, R., Weikert, C., Yuan, J., and Wagner, A. (2011) Adaptation of *Saccharomyces cerevisiae* to saline stress through laboratory evolution. *J Evol Biol* **24**: 1135–1153.
- Díaz Muñoz, G., and Montalvo-Rodríguez, R. (2005) Halophilic black yeast *Hortaea werneckii* in the Cabo Rojo solar salterns: Its first record for this extreme environment in Puerto Rico. *Caribb J Sci* **41**: 360–365.
- Djelal, H., Amrane, A., Lahrer, F., and Martin, G. (2005) Effect of medium osmolarity on the bioproduction of glycerol and ethanol by *Hansenula anomala* growing on glucose and ammonium. *Appl Microbiol Biotechnol* **69**: 341–349.
- Domínguez, J.M., Gong, C.S., and Tsao, G.T. (1997) Production of xylitol from D-xylose by *Debaryomyces hansenii*. *Appl Biochem Biotechnol* **63–65**: 117–127.
- Domínguez-Cuevas, P., González-Pastor, J.-E., Marqués, S., Ramos, J.-L., and de Lorenzo, V. (2006) Transcriptional tradeoff between metabolic and stress-response programs in *Pseudomonas putida* KT2440 cells exposed to toluene. *J Biol Chem* **281**: 11981–11991.
- Duda, V.I., Danilevich, V.N., Suzina, N.E., Shorokhova, A.P., Dmitriev, V.V., Mokhova, O.N., et al. (2004) Changes in the fine structure of microbial cells induced by chaotropic salts. *Microbiology (Russia)* **73**: 341–349.
- Duiff, B.J., Meijer, J.W., Bakker, P.A.H.M., and Schippers, B. (1993) Siderophore-mediated competition for iron and induced resistance in the suppression of *Fusarium* wilt of carnation by fluorescent *Pseudomonas* spp. *Neth J Plant Pathol* **99**: 277–289.
- Egan, M.W., Spratt, D.A., Ng, Y.L., Lam, J.M., Moles, D.R., and Gulabivala, K. (2002) Prevalence of yeasts in saliva and root canals of teeth associated with apical periodontitis. *Int Endod J* **35**: 321–329.
- El Ghachi, M., Bouhss, A., Blanot, D., and Mengin-Lecreulx, D. (2004) The *bacA* gene of *Escherichia coli* encodes an undecaprenyl pyrophosphate phosphatase activity. *J Biol Chem* **279**: 30106–30113.
- Enya, J., Shinohara, H., Yoshida, S., Negishi, T.T.H., Suyama, K., and Tsushima, S. (2007) Culturable leaf-associated bacteria on tomato plants and their potential as biological control agents. *Microb Ecol* **53**: 524–536.
- Escapa, I.F., García, J.L., Bühler, B., Blank, L.M., and Prieto, M.A. (2012) The polyhydroxyalkanoate metabolism controls carbon and energy spillage in *Pseudomonas putida*. *Environ Microbiol* **14**: 1049–1063.
- Eshkol, N., Sendovski, M., Bahalul, M., Katz-Ezov, T., Kashi, Y., and Fishman, A. (2009) Production of 2-phenylethanol from L-phenylalanine by a stress tolerant *Saccharomyces cerevisiae* strain. *J Appl Microbiol* **106**: 534–542.
- Ezeji, T., Milne, C., Price, N., and Blaschek, H. (2010) Achievements and perspectives to overcome the poor solvent resistance in acetone and butanol-producing microorganisms. *Appl Microbiol Biotechnol* **85**: 1697–1712.
- Farkas, I., Hardy, T.A., DePaoli-Roach, A.A., and Roach, P.J. (1990) Isolation of the GSY1 gene encoding yeast glycerol synthase and evidence for the existence of a second gene. *J Biol Chem* **265**: 20879–20886.
- Favero, M.S., Carson, L.A., Bond, W.W., and Petersen, N.J. (1971) *Pseudomonas aeruginosa* – growth in distilled water from hospitals. *Science* **173**: 836–838.

- Ferrer, M., Chernikova, T.N., Yakimov, M.M., Golyshin, P.N., and Timmis, K.N. (2003) Chaperonins govern growth of *Escherichia coli* at low temperatures. *Nat Biotechnol* **21**: 1266–1267.
- Fialho, M.B., Ferreira, L.F.R., Monteiro, R.T.R., and Paschoalati, S.F. (2011) Antimicrobial volatile organic compounds affect morphogenesis-related enzymes in *Guignardia citricarpa*, causal agent of citrus black spot. *Biocontrol Sci Technol* **21**: 797–807.
- Fillet, S., Daniels, C., Pini, C., Krell, T., Duque, E., Bernal, P., et al. (2012) Transcriptional control of the main aromatic hydrocarbon efflux pump in *Pseudomonas*. *Environ Microbiol Rep* **4**: 158–167.
- Fleming, A. (1929) On the antibacterial action of cultures of a *Penicillium*, with special reference to their use in the isolation of *B. influenza*. *Exp Pathol* **10**: 226–236.
- Flipphi, M., Sun, J.B., Robellet, X., Karaffa, L., Fekete, E., Zeng, A.P., et al. (2009) Biodiversity and evolution of primary carbon metabolism in *Aspergillus nidulans* and other *Aspergillus* spp. *Fungal Genet Biol* **46**: S19–S44.
- Freitas, J.S., Silva, E.M., Leal, J., Gras, D.E., Martinez-Rossi, N.M., dos Santos, L.D., et al. (2011) Transcription of the Hsp30, Hsp70, and Hsp90 heat shock protein genes is modulated by the PalA protein in response to acid pH-sensing in the fungus *Aspergillus nidulans*. *Cell Stress Chaperones* **16**: 565–572.
- Gajic, O., Buist, G., Kojic, M., Topisirovic, L., Kuipers, O.P., and Kok, J. (2003) Novel mechanism of bacteriocin secretion and immunity carried out by Lactococcal multidrug resistance proteins. *J Biol Chem* **278**: 34291–34298.
- Garvie, E.I. (1980) Bacterial lactate dehydrogenases. *Microbiol Rev* **44**: 106–139.
- George, H.A., Johnson, J.L., Moore, W.E.C., Holdeman, L.V., and Chen, J.S. (1983) Acetone, isopropanol, and butanol production by *Clostridium beijerinckii* (syn. *Clostridium butylicum*) and *Clostridium aurantibutyricum*. *Appl Environ Microbiol* **45**: 1160–1163.
- Golubev, V.I., and Churkina, L.G. (1993) Intrageneric killing patterns of *Rhodotorula mucilaginosa* mycocins. *Izv Akad Nauk Ser Biol* (4): 550–557.
- Golubev, V.I., Churkina, L.G., and Seregina, S.A. (1996) Intergeneric spectra of action of the *Rhodotorula mucilaginosa* mycocins. *Izv Akad Nauk Ser Biol* (5): 523–529.
- Görner, W., Durchschlag, E., Martinez-Pastor, M.T., Estruch, F., Ammerer, G., Hamilton, B., et al. (1998) Nuclear localization of the C₂H₂ zinc finger protein Msn2p is regulated by stress and protein kinase A activity. *Genes Dev* **12**: 586–597.
- Gostinčar, C., Grube, M., de Hoog, S., Zalar, P., and Gunde-Cimerman, N. (2010) Extremotolerance in fungi: evolution on the edge. *FEMS Microbiol Ecol* **71**: 2–11.
- Gostinčar, C., Gunde-Cimerman, N., and Turk, M. (2012) Genetic resources of extremotolerant fungi: a method for identification of genes conferring stress tolerance. *Biore-sour Technol* **111**: 360–367.
- Goyal, R., Das, S., Arora, A., and Aggarwal, A. (2008) *Rhodotorula mucilaginosa* as a cause of persistent femoral nonunion. *J Postgrad Med* **54**: 25–27.
- Grant, W.D. (2004) Life at low water activity. *Philos Trans R Soc Lond B Biol Sci* **359**: 1249–1266.
- Guillot, A., Gitton, C., Anglade, P., and Mistou, M.-Y. (2003) Proteomic analysis of *Lactococcus lactis*, a lactic acid bacterium. *Proteomics* **3**: 337–354.
- Guillou, V., Plourde-Owobi, L., Parrou, J.L., Goma, G., and François, J. (2004) Role of reserve carbohydrates in the growth dynamics of *Saccharomyces cerevisiae*. *FEMS Yeast Res* **4**: 773–787.
- Gulati, A., Sharma, N., Vyas, P., Sood, S., Rahi, P., Pathania, V., et al. (2010) Organic acid production and plant growth promotion as a function of phosphate solubilization by *Acinetobacter rhizosphaerae* strain BIHB 723 isolated from the cold deserts of the trans-Himalayas. *Arch Microbiol* **192**: 975–983.
- Gunde-Cimerman, N., Zalar, P., de Hoog, S., and Plemenitas, A. (2000) Hypersaline waters in salterns – natural ecological niches for halophilic black yeasts. *FEMS Microbiol Ecol* **32**: 235–240.
- Hallsworth, J.E. (1998) Ethanol-induced water stress in yeast. *J Ferment Bioeng* **85**: 125–137.
- Hallsworth, J.E., and Magan, N. (1994a) Effect of carbohydrate type and concentration on polyols and trehalose in conidia of three entomopathogenic fungi. *Microbiology* **140**: 2705–2713.
- Hallsworth, J.E., and Magan, N. (1994b) Improved biological control by changing polyols/trehalose in conidia of entomopathogens. In *Brighton Crop Protection Council – Pests And Diseases 1994*. McKim, F.M. (ed.). Farnham, Surrey, UK: British Crop Protection Council, pp. 1091–1096.
- Hallsworth, J.E., and Magan, N. (1994c) Effects of KCl concentration on accumulation of acyclic sugar alcohols and trehalose in conidia of three entomopathogenic fungi. *Lett Appl Microbiol* **18**: 8–11.
- Hallsworth, J.E., and Magan, N. (1995) Manipulation of intracellular glycerol and erythritol enhances germination of conidia at low water availability. *Microbiology* **141**: 1109–1115.
- Hallsworth, J.E., and Nomura, Y. (1999) A simple method to determine the water activity of ethanol-containing samples. *Biotechnol Bioeng* **62**: 242–245.
- Hallsworth, J.E., Nomura, Y., and Iwahara, M. (1998) Ethanol-induced water stress and fungal growth. *J Ferment Bioeng* **86**: 451–456.
- Hallsworth, J.E., Heim, S., and Timmis, K.N. (2003a) Chaotropic solutes cause water stress in *Pseudomonas putida*. *Environ Microbiol* **5**: 1270–1280.
- Hallsworth, J.E., Prior, B.A., Nomura, Y., Iwahara, M., and Timmis, K.N. (2003b) Compatible solutes protect against chaotrope (ethanol)-induced, nonosmotic water stress. *Appl Environ Microbiol* **69**: 7032–7034.
- Hallsworth, J.E., Yakimov, M.M., Golyshin, P.N., Gillion, J.L.M., D'Auria, G., Alves, F.D.L., et al. (2007) Limits of life in MgCl₂-containing environments: chaotropicity defines the window. *Environ Microbiol* **9**: 801–813.
- Hartmans, S., Smits, J.P., van der Werf, M.J., Volkerling, F., and de Bont, J.A.M. (1989) Metabolism of styrene oxide and 2-phenylethanol in the styrene-degrading *Xanthobacter* strain 124X. *Appl Environ Microbiol* **55**: 2850–2855.
- Hassan, M.N., Afghan, S., and Hafeez, F.Y. (2011) Biological control of red rot in sugarcane by native pyoluteorin-producing *Pseudomonas putida* strain NH-50 under field

- conditions and its potential modes of action. *Pest Manag Sci* **67**: 1147–1154.
- Hellström, A.M., Almgren, A., Carlsson, N.-G., Svanberg, U., and Andlid, T.A. (2012) Degradation of phytate by *Pichia kudriavzevii* TY13 and *Hanseniaspora guilliermondii* TY14 in Tanzanian togwa. *Int J Food Microbiol* **153**: 73–77.
- Hentzer, M., Teitzel, G.M., Balzer, G.J., Heydorn, A., Molin, S., Givskov, M., *et al.* (2001) Alginate overproduction affects *Pseudomonas aeruginosa* biofilm structure and function. *J Bacteriol* **183**: 5395–5401.
- Herrero, M., Ibáñez, E., Cifuentes, A., Reglero, G., and Santoyo, S. (2006) *Dunaliella salina* microalga pressurized liquid extracts as potential antimicrobials. *J Food Prot* **69**: 2471–2477.
- Hill, T.A. (1977) *The Biology of Weeds*. London, UK: E. Arnold.
- Hirasawa, T., Yoshikawa, K., Nakakura, Y., Nagahisa, K., Furusawa, C., Katakura, Y., *et al.* (2007) Identification of target genes conferring ethanol stress tolerance to *Saccharomyces cerevisiae* based on DNA microarray data analysis. *J Bacteriol* **131**: 34–44.
- Hirsch, P. (1986) Microbial life at extremely low nutrient levels. *Adv Space Res* **6**: 287–298.
- Hocking, A.D. (1993) Responses of xerophilic fungi to changes in water activity. In *Stress Tolerance of Fungi*. Jennings, D.H. (ed.). New York, USA: Marcel Dekker, pp. 233–256.
- Hoffland, E., Kuyper, T.W., Wallander, H., Plassard, C., Gorbushina, A.A., Haselwandter, K., *et al.* (2004) The role of fungi in weathering. *Front Ecol Environ* **2**: 258–264.
- Hong, J., Tamaki, H., Akiba, S., Yamamoto, K., and Kumagai, H. (2001) Cloning of a gene encoding a highly stable endo- β -1,4-glucanase from *Aspergillus niger* and its expression in yeast. *J Biosci Bioeng* **92**: 434–441.
- Horinouchi, T., Tamaoka, K., Furusawa, C., Ono, N., Suzuki, S., Hirasawa, T., *et al.* (2010) Transcriptome analysis of parallel-evolved *Escherichia coli* strains under ethanol stress. *BMC Genomics* **11**: 579.
- Hunter, P.J., Hand, P., Pink, D., Whipps, J.M., and Bending, G.D. (2010) Both leaf properties and microbe-microbe interactions influence within-species variation in bacterial population diversity and structure in the lettuce (*Lactuca* species) phyllosphere. *Appl Environ Microbiol* **76**: 8117–8125.
- Inderjit, Evans, H., Crocoll, C., Bajpai, D., Kaur, R., Feng, Y.L., *et al.* (2011) Volatile chemicals from leaf litter are associated with invasiveness of a Neotropical weed in Asia. *Ecology* **92**: 316–324.
- Ingram, L.O., and Buttke, T.M. (1984) Effects of alcohols on microorganisms. *Adv Microb Physiol* **25**: 253–300.
- Izgü, F., Altınbay, D., and Sertkaya, A. (2005) Enzymic activity of the K5-type yeast killer toxin and its characterization. *Biosci Biotechnol Biochem* **69**: 2200–2206.
- Jahid, I.K., Silva, A.J., and Benitez, J.A. (2006) Polyphosphate stores enhance the ability of *Vibrio cholerae* to overcome environmental stresses in a low-phosphate environment. *Appl Environ Microbiol* **72**: 7043–7049.
- Jang, M.-H., Ha, K., Joo, G.-J., and Takamura, N. (2003) Toxin production of cyanobacteria is increased by exposure to zooplankton. *Freshw Biol* **48**: 1540–1550.
- Jiménez, J.I., Miñambres, B., García, J.L., and Díaz, E. (2002) Genomic analysis of the aromatic catabolic pathways from *Pseudomonas putida* KT2440. *Environ Microbiol* **4**: 824–841.
- Jorge, R.M.F., and Livingston, A.G. (1999) A novel method for characterisation of microbial growth kinetics on volatile organic compounds. *Appl Microbiol Biotechnol* **52**: 174–178.
- Jüttner, F. (2005) Evidence that polyunsaturated aldehydes of diatoms are repellents for pelagic crustacean grazers. *Aquat Ecol* **39**: 271–282.
- Jüttner, F., Watson, S.B., von Elert, E., and Koster, O. (2010) β -Cyclocitral, a grazer defence signal unique to the cyanobacterium *Microcystis*. *J Chem Ecol* **36**: 1387–1397.
- Kachalkin, A.V., and Yurkov, A.M. (2012) Yeast communities in Sphagnum phyllosphere along the temperature-moisture ecocline in the boreal forest-swamp ecosystem and description of *Candida sphagnicola* sp. nov. *Antonie Van Leeuwenhoek* **102**: 29–43.
- Kadouri, D., Jurkevitch, E., Okon, Y., and Castro-Sowinski, S. (2005) Ecological and agricultural significance of bacterial polyhydroxyalkanoates. *Crit Rev Microbiol* **31**: 55–67.
- Kaino, T., and Takagi, H. (2008) Gene expression profiles and intracellular contents of stress protectants in *Saccharomyces cerevisiae* under ethanol and sorbitol stresses. *Appl Microbiol Biotechnol* **79**: 273–283.
- Kanekar, S.P., Kanekar, P.P., Sarnaik, S.S., Gujrathi, N.P., Shede, P.N., Kedargol, M.R., and Reardon, K.F. (2009) Bioremediation of nitroexplosive wastewater by an yeast isolate *Pichia sydowiorum* MCM Y-3 in fixed film bioreactor. *J Ind Microbiol Biotechnol* **36**: 253–260.
- Karlin, S., Mrázek, J., Campbell, A., and Kaiser, D. (2001) Characterizations of highly expressed genes of four fast-growing bacteria. *J Bacteriol* **183**: 5025–5040.
- Kashangura, C., Hallsworth, J.E., and Mswaka, A.Y. (2006) Phenotypic diversity amongst strains of *Pleurotus sajor-caju*: implications for cultivation in arid environments. *Mycol Res* **110**: 312–317.
- Kazda, J., and Falkinham, J.O., III (2009) Mycobacteria in sphagnum, peats and potting soils. In *The Ecology of Mycobacteria: Impact on Animal's and Human's Health*. Kazda, J., Pavlik, I., Falkinham, J.O., III, and Hruska, K. (eds). Heidelberg: Springer, pp. 89–95.
- Kets, E.P.W., Galinski, E.A., de Wit, M., de Bont, J.A.M., and Heipieper, H.J. (1996) Mannitol, a novel bacterial compatible solute in *Pseudomonas putida* S12. *J Bacteriol* **178**: 6665–6670.
- Khemakhem, H., Elloumi, J., Moussa, M., Aleya, L., and Ayadi, H. (2010) The concept of ecological succession applied to phytoplankton over four consecutive years in five ponds featuring a salinity gradient. *Estuar Coast Shelf Sci* **88**: 33–44.
- Kiran, G.S., Hema, T.A., Gandhimathi, R., Selvin, J., Thomas, T.A., Rajeetha Ravji, T., *et al.* (2009) Optimization and production of a biosurfactant from the sponge-associated marine fungus *Aspergillus ustus* MSF3. *Colloids Surf B Biointerfaces* **73**: 250–256.
- Klappenbach, J.A., Dunbar, J.M., and Schmidt, T.M. (2000) rRNA operon copy number reflects ecological strategies of bacteria. *Appl Environ Microbiol* **66**: 1328–1333.

- van der Klei, I.J., Harder, W., and Veenhuis, M. (1991) Biosynthesis and assembly of alcohol oxidase, a peroxisomal matrix protein in methylotrophic yeasts: a review. *Yeast* **7**: 195–209.
- Klinke, T., Kneist, S., de Soet, J.J., Kuhlisch, E., Mauersberger, S., Förster, A., et al. (2009) Acid production by oral strains of *Candida albicans* and Lactobacilli. *Caries Res* **43**: 83–91.
- Koch, A.L. (2001) Oligotrophs versus copiotrophs. *Bioessays* **23**: 657–661.
- Kolaczowski, M., van der Rest, M., Cybularz-Kolaczowska, A., Soumilion, J.-P., Konings, W.N., and Goffeau, A. (1996) Anticancer drugs, ionophoric peptides, and steroids as substrates of the yeast multidrug transporter Pdr5p. *J Biol Chem* **271**: 31543–31548.
- Koornneef, M., and Meinke, D. (2010) The development of *Arabidopsis* as a model plant. *Plant J* **61**: 909–921.
- Kotyk, A., Lapathitis, G., and Krenková, S. (1999) Glucose- and K⁺-induced acidification in different yeast species. *Folia Microbiol (Praha)* **44**: 295–298.
- Krallish, I., Gonta, S., Savenkova, L., Bergauer, P., and Margesin, R. (2006) Phenol degradation by immobilized cold-adapted yeast strains of *Cryptococcus terreus* and *Rhodotorula creatinivora*. *Extremophiles* **10**: 441–449.
- Krappmann, S., and Braus, G.H. (2005) Nitrogen metabolism of *Aspergillus* and its role in pathogenicity. *Med Mycol* **43** (Suppl. 1): S31–S40.
- Kruijt, M., Tran, H., and Raaijmakers, J.M. (2009) Functional, genetic and chemical characterization of biosurfactants produced by plant growth-promoting *Pseudomonas putida* 267. *J Appl Microbiol* **107**: 546–556.
- Kushner, D.J. (ed.) (1978) Life at high salt and solute concentrations: halophilic bacteria. In *Microbial Life in Extreme Environments*. London, UK: Academic Press, pp. 318–368.
- Lahav, R., Fareleira, P., Nejdat, A., and Abeliovich, A. (2002) The identification and characterization of osmotolerant yeast isolates from chemical wastewater evaporation ponds. *Microb Ecol* **43**: 388–396.
- Lahav, R., Nejdat, A., and Abeliovich, A. (2004) Alterations in protein synthesis and levels of heat shock 70 proteins in response to salt stress of the halotolerant yeast *Rhodotorula mucilaginosa*. *Antonie Van Leeuwenhoek* **85**: 259–269.
- Lambou, K., Lamarre, C., Beau, R., Dufour, N., and Latge, J.P. (2010) Functional analysis of the superoxide dismutase family in *Aspergillus fumigatus*. *Mol Microbiol* **75**: 910–923.
- Langaee, T.Y., Gagnon, L., and Huletsky, A. (2000) Inactivation of the *ampD* Gene in *Pseudomonas aeruginosa* leads to moderate-basal-level and hyperinducible AmpC β -lactamase expression. *Antimicrob Agents Chemother* **44**: 583–589.
- Le Bihan, T., Martin, S.F., Chirnside, E.S., van Ooijen, G., Barrios-Llerena, M.E., O'Neill, J.S., et al. (2011) Shotgun proteomic analysis of the unicellular alga *Ostreococcus tauri*. *J Proteomics* **74**: 2060–2070.
- Leão, P.N., Ramos, V., Vale, M., Machado, J.P., and Vasconcelos, V.M. (2012) Microbial community changes elicited by exposure to cyanobacterial allelochemicals. *Microb Ecol* **63**: 85–95.
- Lee, G.D., Shin, S.Y., Maeng, C.-Y., Jin, Z.Z., Kim, K.L., and Hahn, K.-S. (1999) Isolation and characterization of a novel antifungal peptide from *Aspergillus niger*. *Biochem Biophys Res Commun* **263**: 646–651.
- Lee, Y.J., Choi, Y.R., Lee, S.Y., Park, J.T., Shim, J.H., Park, K.H., et al. (2011) Screening wild yeast strains for alcohol fermentation from various Fruits. *Mycobiology* **39**: 33–39.
- Leong, S.L., Pettersson, O.V., Rice, T., Hocking, A.D., and Schnürer, J. (2011) The extreme xerophilic mould *Xeromyces bisporus* – growth and competition at various water activities. *Int J Food Microbiol* **145**: 57–63.
- Lesuisse, E., Blaiseau, P.L., Dancis, A., and Camadro, J.M. (2001) Siderophore uptake and use by the yeast *Saccharomyces cerevisiae*. *Microbiology* **147**: 289–298.
- Levine, D.W., and Cooney, C.L. (1973) Isolation and characterization of a thermotolerant methanol-utilizing yeast. *Appl Microbiol* **26**: 982–990.
- Liberman, L.M., Sozzani, R., and Benfey, P.N. (2012) Integrative systems biology: an attempt to describe a simple weed. *Curr Opin Plant Biol* **15**: 162–167.
- Libkind, D., and Sampaio, J.P. (2010) Rhodotorula. In *Molecular Detection of Foodborne Pathogens*. Liu, D. (ed.). Florida, USA: CRC Press, pp. 603–618.
- Lifshits, M., and Carmeli, S. (2012) Metabolites of *Microcystis aeruginosa* bloom material from Lake Kinneret, Israel. *J Nat Prod* **75**: 209–219.
- Lillie, S.H., and Pringle, J.R. (1980) Reserve carbohydrate metabolism in *Saccharomyces cerevisiae*: responses to nutrient limitation. *J Bacteriol* **143**: 1384–1394.
- Lisichkina, G.A., Bab'eva, I.P., and Dlu, S. (2003) Alkalitolerant yeasts from natural biotopes. *Mikrobiologia* **72**: 695–698.
- Liu, I.-C., Whang, L.-M., Ren, W.-J., and Lin, P.-Y. (2011) The effect of pH on the production of biohydrogen by clostridia: thermodynamic and metabolic considerations. *Int J Hydrogen Energy* **36**: 439–449.
- Liu, J., and Nikaido, H. (1999) A mutant of *Mycobacterium smegmatis* defective in the biosynthesis of mycolic acids accumulates meromycolates. *Proc Natl Acad Sci USA* **96**: 4011–4016.
- Lo, Y.-C., Lu, W.-C., Chen, C.-Y., Chen, W.-M., and Chang, J.-S. (2010) Characterization and high-level production of xylanase from an indigenous cellulolytic bacterium *Acinetobacter junii* F6-02 from southern Taiwan soil. *Biochem Eng J* **53**: 77–84.
- Lo Nostro, P., Ninham, B.W., Lo Nostro, A., Pesavento, G., Fratoni, L., and Baglioni, P. (2005) Specific ion effects on the growth rates of *Staphylococcus aureus* and *Pseudomonas aeruginosa*. *Phys Biol* **2**: 1–7.
- Loaces, I., Ferrando, L., and Scavino, A.F. (2011) Dynamics, diversity and function of endophytic siderophore-producing bacteria in rice. *Microb Ecol* **61**: 606–618.
- Lobasso, S., Lopalco, P., Vitale, R., Saponetti, M.S., Capitanio, G., Mangini, V., et al. (2012) The light-activated proton pump Bop I of the archaeon *Haloquadratum walsbyi*. *Photochem Photobiol* **88**: 690–700.
- Long, H.C. (1932) *Weeds of Grass Land*. London, UK: H.M.S.O..
- Louvel, B., Cébron, A., and Leyval, C. (2011) Root exudates affect phenanthrene biodegradation, bacterial community and functional gene expression in sand microcosms. *Int Biodeter Biodegr* **65**: 947–953.

- Lowe, D.P., and Arendt, E.K. (2004) The use and effects of lactic acid bacteria in malting and brewing with their relationships to antifungal activity, mycotoxins and gushing: a review. *J Inst Brew* **110**: 163–180.
- Ma, M., and Liu, L.Z. (2010) Quantitative transcription dynamic analysis reveals candidate genes and key regulators for ethanol tolerance in *Saccharomyces cerevisiae*. *BMC Microbiol* **10**: 169.
- McArthur, J.V. (2006) *Microbial Ecology: An Evolutionary Approach*. Oxford, UK: Elsevier.
- McCamick, E.M., Gomase, V.S., Timson, D.J., McGenity, T.J., and Hallsworth, J.E. (2009) Water-hydrophobic compound interactions with the microbial cell. In *Handbook of Hydrocarbon and Lipid Microbiology – Hydrocarbons, Oils and Lipids: Diversity, Properties and Formation*, Vol. 2. Timmis, K.N. (ed.). New York, USA: Springer, pp. 1451–1466.
- McDonald, I.R., and Murrell, J.C. (1997) The methanol dehydrogenase structural gene *mxhF* and its use as a functional gene probe for methanotrophs and methylotrophs. *Appl Environ Microbiol* **63**: 3218–3224.
- McDonnell, G., and Russell, A.D. (1999) Antiseptics and disinfectants: activity, action, and resistance. *Clin Microbiol Rev* **12**: 147–179.
- Mack, R.N., Simberloff, D., Lonsdale, W.M., Evans, H., Clout, M., and Bazzaz, F. (2000) Biotic invasions: causes, epidemiology, global consequences, and control. *Ecol Appl* **10**: 689–710.
- McNerney, R., Mallard, K., Okolo, P.I., and Turner, C. (2012) Production of volatile organic compounds by mycobacteria. *FEMS Microbiol Lett* **328**: 150–156.
- Magan, N., Hope, R., Cairns, V., and Aldred, D. (2003) Post-harvest fungal ecology: impact of fungal growth and mycotoxin accumulation in stored grain. *Eur J Plant Pathol* **109**: 723–730.
- Mansure, J.J.C., Panek, A.D., Crowe, L.M., and Crowe, J.H. (1994) Trehalose inhibits ethanol effects on intact yeast cells and liposomes. *Biochim Biophys Acta* **1191**: 309–316.
- Margesin, R., Labbe, D., Schinner, F., Greer, C.W., and Whyte, L.G. (2003) Characterization of hydrocarbon-degrading microbial populations in contaminated and pristine alpine soils. *Appl Environ Microbiol* **69**: 3085–3092.
- Matz, C., Bergfeld, T., Rice, S.A., and Kjelleberg, S. (2004) Microcolonies, quorum sensing and cytotoxicity determine the survival of *Pseudomonas aeruginosa* biofilms exposed to protozoan grazing. *Environ Microbiol* **6**: 218–226.
- Mavrodi, D.V., Blankenfeldt, W., and Thomashow, L.S. (2006) Phenazine compounds in fluorescent *Pseudomonas* spp. biosynthesis and regulation. *Annu Rev Phytopathol* **44**: 417–445.
- Meagher, M.M., Tao, B.Y., Chow, J.M., and Reilly, P.J. (1988) Kinetics and subsite mapping of a D-xylobiose- and D-xylose-producing *Aspergillus niger* endo-(1→4)-β-D-xylanase. *Carbohydr Res* **173**: 273–283.
- Meksawat, S., and Pornprom, T. (2010) Allelopathic effect of itchgrass (*Rottboellia cochinchinensis*) on seed germination and plant growth. *Weed Biol Manag* **10**: 16–24.
- Middelhoven, W.J., de Waard, M.A., and Mulder, E.G. (1969) The ferrous ion as the cofactor of arginase *in vivo*. II. Experiments on the replacement of ferrous ions in native yeast arginase by other cations *in vivo*. *Biochim Biophys Acta* **191**: 122–129.
- Minerdi, D., Bossi, S., Gullino, M.L., and Garibaldi, A. (2009) Volatile organic compounds: a potential direct long-distance mechanism for antagonistic action of *Fusarium oxysporum* strain MSA 35. *Environ Microbiol* **11**: 844–854.
- Minnikin, D.E., Abdolrahimzadeh, H., and Baddiley, J. (1974) Replacement of acidic phosphates by acidic glycolipids in *Pseudomonas diminuta*. *Nature* **249**: 268–269.
- Mlouka, A., Comte, K., Castets, A.M., Bouchier, C., and de Marsac, N. (2004) The gas vesicle gene cluster from *Microcystis aeruginosa* and DNA rearrangements that lead to loss of cell buoyancy. *J Bacteriol* **186**: 2355–2365.
- Moliné, M., Flores, M.R., Libkind, D., Diéguez, M.D.C., Farías, M.E., and van Broock, M. (2010) Photoprotection by carotenoid pigments in the yeast *Rhodotorula mucilaginosa*: the role of torularhodin. *Photochem Photobiol Sci* **9**: 1145–1151.
- Molly, K., DeSmet, I., Nollet, L., Vande Woestyne, M., and Verstraete, W. (1996) Effect of lactobacilli on the ecology of the gastro-intestinal microbiota cultured in the SHIME reactor. *Microb Ecol Health* **9**: 79–89.
- Mongodin, E.F., Nelson, K.E., Daugherty, S., DeBoy, R.T., Wister, J., Khouri, H., et al. (2005) The genome of *Salinibacter ruber*: convergence and gene exchange among hyperhalophilic bacteria and archaea. *Proc Natl Acad Sci USA* **102**: 18147–18152.
- Morales, P., Fernández-García, E., and Nuñez, M. (2005) Volatile compounds produced in cheese by *Pseudomonas* strains of dairy origin belonging to six different species. *J Agric Food Chem* **53**: 6835–6843.
- Morohoshi, T., Oseki, K., and Ikeda, T. (2011) Exopolysaccharide production is influenced by sugars, N-acylhomoserine, lactone, and transcriptional regulators RcsA and RcsB, but does not affect pathogenicity in the plant pathogen *Pantoea ananatis*. *Biosci Biotechnol Biochem* **75**: 997–999.
- Moskvina, E., Schüller, C., Maurer, C.T.C., Mager, W.H., and Ruis, H. (1998) A search in the genome of *Saccharomyces cerevisiae* for genes regulated via stress response elements. *Yeast* **14**: 1041–1050.
- Muñoz, J., Mudge, S.M., and Sandoval, A. (2004) Effects of ionic strength on the production of short chain volatile hydrocarbons by *Dunaliella salina* (Teodoresco). *Chemosphere* **54**: 1267–1271.
- Nair, S., and Finkel, S.E. (2004) Dps protects cells against multiple stresses during stationary phase. *J Bacteriol* **186**: 4192–4198.
- Narihiro, T., and Sekiguchi, Y. (2007) Microbial communities in anaerobic digestion processes for waste and wastewater treatment: a microbiological update. *Curr Opin Biotechnol* **18**: 273–278.
- Naumov, G.I., Naumova, E.S., and Schnürer, J. (2001) Genetic characterization of the nonconventional yeast *Hansenula anomala*. *Res Microbiol* **152**: 551–562.
- Newton, R.J., Kent, A.D., Triplett, E.W., and McMahon, K.D. (2006) Microbial community dynamics in a humic lake: differential persistence of common freshwater phylotypes. *Environ Microbiol* **8**: 956–970.

- Nishino, K., and Yamaguchi, A. (2001) Analysis of a complete library of putative drug transporter genes in *Escherichia coli*. *J Bacteriol* **183**: 5803–5812.
- Nisiotou, A.A., Spiropoulos, A.E., and Nychas, G.E. (2007) Yeast community structures and dynamics in healthy and *Botrytis*-affected grape must fermentations. *Appl Environ Microbiol* **73**: 6705–6713.
- Nisiotou, A.A., Chorianopoulos, N., Nychas, G.-J.E., and Panagou, E.Z. (2010) Yeast heterogeneity during spontaneous fermentation of black Conservolea olives in different brine solutions. *J Appl Microbiol* **108**: 396–405.
- Nitschke, M., Costa, S.G.V.A.O., and Contiero, J. (2011) Rhamnolipids and PHAs: recent reports on *Pseudomonas*-derived molecules of increasing industrial interest. *Process Biochem* **46**: 621–630.
- Nomura, Y., Hallsworth, J.E., Iwahara, M., Tanaka, T., and Ishizaki, A. (1998) Rapid and effective production of L-lactate from xylose using electrodialysis culture-associated product separation. *World J Microb Biot* **14**: 911–916.
- Nordström, K., and Dasgupta, S. (2006) Copy-number control of the *Escherichia coli* chromosome: a plasmidologist's view. *EMBO Rep* **7**: 484–489.
- Nowak-Thompson, B., Chaney, N., Wing, J.S., Gould, S.J., and Loper, J.E. (1999) Characterization of the pyoluteorin biosynthetic gene cluster of *Pseudomonas fluorescens* Pf-5. *J Bacteriol* **181**: 2166–2174.
- Nunes, F.M.N., Veloso, M.C.C., Pereira, P.A.P., and de Andrade, J.B. (2005) Gas-phase ozonolysis of the monoterpene (S)-(+)-carvone, (R)-(–)-carvone, (–)-carveol, geraniol and citral. *Atmos Environ* **39**: 7715–7730.
- Oatham, M.P. (1997) Water relations of *Zygophyllum hamense*, *Heliotropium kotschy* and *Panicum turgidum* and their response to rangeland management techniques in Abu Dhabi. *J Arid Environ* **35**: 95–110.
- Ogino, A., Koshikawa, H., Nakahara, T., and Uchiyama, H. (2001) Succession of microbial communities during a biostimulation process as evaluated by DGGE and clone library analyses. *J Appl Microbiol* **91**: 625–635.
- Olstorpe, M., and Passoth, M. (2011) *Pichia anomala* in grain preservation. *Antonie Van Leeuwenhoek* **99**: 57–62.
- Oren, A. (1999) Bioenergetic aspects of halophilism. *Microbiol Mol Biol Rev* **63**: 334–338.
- Oren, A., Heldal, M., Norland, S., and Galinski, E.A. (2002) Intracellular ion and organic solute concentrations of the extremely halophilic bacterium *Salinibacter ruber*. *Extremophiles* **6**: 491–498.
- Ozaki, K., Ohta, A., Iwata, C., Horikawa, A., Tsuji, K., Ito, E., et al. (2008) Lysis of cyanobacteria with volatile organic compounds. *Chemosphere* **71**: 1531–1538.
- Pacheco, C.C., Passos, J.F., Moradas-Ferreira, P., and De Marco, P. (2003) Strain PM2, a novel methylotrophic fluorescent *Pseudomonas* sp. *FEMS Microbiol Lett* **227**: 279–285.
- Pan, F., Yang, Q., Zhang, Y., Zhang, S., and Yang, M. (2004) Biodegradation of polycyclic aromatic hydrocarbons by *Pichia anomala*. *Biotechnol Lett* **26**: 803–806.
- Park, W., Jeon, C.O., Cadillo, H., DeRito, C., and Madsen, E.L. (2004) Survival of naphthalene-degrading *Pseudomonas putida* NCIB 9816-4 in naphthalene-amended soils: toxicity of naphthalene and its metabolites. *Appl Microbiol Biotechnol* **64**: 429–435.
- Park, H.C., Bae, Y.U., Cho, S.D., Kim, S.A., Moon, J.Y., Ha, K.C., et al. (2007) Toluene-induced accumulation of trehalose by *Pseudomonas* sp. BCNU 106 through expression of otsA and otsB homologues. *Lett Appl Microbiol* **1**: 50–55.
- Parkinson, N., Bryant, R., Bew, J., and Elphinstone, J. (2011) Rapid phylogenetic identification of members of the *Pseudomonas syringae* species complex using the *rpoD* locus. *Plant Pathol* **60**: 338–344.
- Parret, A.H.A., Schoofs, G., Proost, P., and Mot, D.M. (2003) Plant lectin-like bacteriocin from a rhizosphere-colonizing *Pseudomonas* isolate. *J Bacteriol* **185**: 897–908.
- Parrou, J.L., Teste, M.A., and François, J. (1997) Effects of various types of stress on the metabolism of reserve carbohydrates in *Saccharomyces cerevisiae*: genetic evidence for a stress-induced recycling of glycogen and trehalose. *Microbiology* **143**: 1891–1900.
- Passoth, V., Fredlund, E., Druvefors, U., and Schnürer, J. (2006) Biotechnology, physiology and genetics of the yeast *Pichia anomala*. *FEMS Yeast Res* **6**: 3–13.
- Paul, D., Pandey, G., Meier, C., van der Meer, J.R., and Jain, R.K. (2006) Bacterial community structure of a pesticide-contaminated site and assessment of changes induced in community structure during bioremediation. *FEMS Microbiol Ecol* **57**: 116–127.
- Paz, Y., Shimoni, E., Weiss, M., and Pick, U. (2007) Effects of iron deficiency on iron binding and internalization into acidic vacuoles in *Dunaliella salina*. *Plant Physiol* **144**: 1407–1415.
- Pecoraro, V., Zerulla, K., Lange, C., and Soppa, J. (2011) Quantification of ploidy in proteobacteria revealed the existence of monoploid, (mero-)oligoploid and polyploid species. *PLoS ONE* **6**: e16392.
- Peltomaa, E., and Ojala, A. (2010) Size-related photosynthesis of algae in a strongly stratified humic lake. *J Plankton Res* **32**: 341–355.
- Perniola, R., Faneschi, M.L., Manso, E., Pizzolante, M., Rizzo, A., Sticchi Damiani, A., et al. (2006) *Rhodotorula mucilaginosa* outbreak in neonatal intensive care unit: microbiological features, clinical presentation, and analysis of related variables. *Eur J Clin Microbiol Infect Dis* **25**: 193–196.
- Pfeiffer, T., Schuster, S., and Bonhoeffer, S. (2001) Cooperation and competition in the evolution of ATP-producing pathways. *Science* **292**: 504–507.
- de Pina, C.G., and Hogg, T.A. (1999) Microbial and chemical changes during the spontaneous ensilage of grape pomace. *J Appl Microbiol* **86**: 777–784.
- Piškur, J., Rozpędowska, E., Polakova, S., Merico, A., and Compagno, C. (2006) How did *Saccharomyces* evolve to become a good brewer? *Trends Genet* **22**: 183–186.
- Pitt, J.I. (1975) Xerophilic fungi and the spoilage of foods of plant origin. In *Water Relations of Foods*, 1st edn. Duckworth, R.B. (ed.). New York, USA: Academic Press, pp. 273–307.
- Pitt, J.I., and Christian, J.H.B. (1968) Water relations of xerophilic fungi isolated from prunes. *Appl Microbiol* **16**: 1853–1858.

- Pitt, J.I., and Hocking, A.D. (1977) Influence of solute and hydrogen-ion concentration on water relations of some xerophilic fungi. *J Gen Microbiol* **101**: 35–40.
- Poblete-Castro, I., Becker, J., Dohnt, J., dos Santos, V.M., and Wittman, C. (2012) Industrial biotechnology of *Pseudomonas putida* and other related species. *Appl Microbiol Biotechnol* **93**: 2279–2290.
- Pollegioni, L., Piubelli, L., Sacchi, S., Pilone, M.S., and Molla, G. (2007) Physiological functions of D-amino acid oxidases: from yeast to humans. *Cell Mol Life Sci* **64**: 1373–1394.
- Praphailong, W., and Fleet, G.H. (1997) The effect of pH, sodium chloride, sucrose, sorbate and benzoate on the growth of food spoilage yeasts. *Food Microbiol* **14**: 459–468.
- Pretorius, I.S. (2000) Tailoring wine yeast for the new millennium: novel approaches to the ancient art of winemaking. *Yeast* **16**: 675–729.
- Puchałka, J., Oberhardt, M.A., Godinho, M., Bielecka, A., Regenhardt, D., Timmis, K.N., et al. (2008) Genome-scale reconstruction and analysis of the *Pseudomonas putida* KT2440 metabolic network facilitates applications in biotechnology. *PLoS Comput Biol* **4**: e1000210.
- Purvis, J.E., Yomano, L.P., and Ingram, L.O. (2005) Enhanced trehalose production improves growth of *Escherichia coli* under osmotic stress. *Appl Environ Microbiol* **71**: 3761–3769.
- Ragaert, P., Devlieghere, F., Loos, S., Dewulf, J., van Langenhove, H., Foubert, I., et al. (2006) Role of yeast proliferation in the quality degradation of strawberries during refrigerated storage. *Int J Food Microbiol* **108**: 42–50.
- Rainieri, S., Zambonelli, C., Hallsworth, J.E., Pulvirenti, A., and Giudici, P. (1999) *Saccharomyces uvarum*, a distinct group within *Saccharomyces sensu stricto*. *FEMS Microbiol Lett* **177**: 177–185.
- Rajala, T., Peltoniemi, M., Hantula, J., Mäkipää, R., and Penanen, T. (2011) RNA reveals a succession of active fungi during the decay of Norway spruce logs. *Fungal Ecol* **4**: 437–448.
- Ralphy, M.H. (2002) Ecological relationships between poisonous plants and rangeland condition: a review. *J Range Manag* **55**: 285–290.
- Ramos, J.L., Duque, E., Gallegos, M.-T., Godoy, P., Ramos-González, M.I., Rojas, A., et al. (2002) Mechanisms of solvent tolerance in Gram-negative bacteria. *Annu Rev Microbiol* **56**: 743–768.
- Randazzo, C.L., Pitino, I., Ribbera, A., and Caggia, C. (2010) Pecorino Crotonese cheese: study of bacterial population and flavour compounds. *Food Microbiol* **27**: 363–374.
- Rao, D., Webb, J.S., and Kjelleberg, S. (2005) Competitive interactions in mixed-species biofilms containing the marine bacterium *Pseudoalteromonas tunicate*. *Appl Environ Microbiol* **71**: 1729–1736.
- Raspor, P., Milek, D.M., Polanc, J., Možina, S.S., and Čadež, N. (2006) Yeasts isolated from three varieties of grapes cultivated in different locations of the Dolenjska vine-growing region, Slovenia. *Int J Food Microbiol* **109**: 97–102.
- Reis, R.S., Pereira, A.G., Neves, B.C., and Freire, D.M.G. (2011) Gene regulation of rhamnolipid production in *Pseudomonas aeruginosa* – a review. *Bioresour Technol* **102**: 6377–6384.
- Rengefors, K., Pålsson, C., Hansson, L.-A., and Heiberg, L. (2008) Cell lysis of competitors and osmotrophy enhance growth of the bloom-forming alga *Gonyostomum semen*. *Aquat Microb Ecol* **51**: 87–96.
- Renouf, V., Claisse, O., and Lonvaud-Funel, A. (2005) Understanding the microbial ecosystem on the grape berry surface through numeration and identification of yeast and bacteria. *Aust J Grape Wine Res* **11**: 316–327.
- Renshaw, J.C., Robson, G.D., Trinci, A.P.J., Wiebe, M.G., Livens, F.R., Collison, D., et al. (2002) Fungal siderophores: structures, functions and applications. *Mycol Res* **106**: 1123–1142.
- Restaino, L., Bills, S., Tscherneff, K., and Lenovich, L.M. (1983) Growth-characteristics of *Saccharomyces rouxii* isolated from chocolate syrup. *Appl Environ Microbiol* **45**: 1614–1621.
- Reynisson, E., Lauzon, H.L., Magnússon, H., Jónsdóttir, R., Ólafsdóttir, G., Marteinson, V., et al. (2009) Bacterial composition and succession during storage of North-Atlantic cod (*Gadus morhua*) at superchilled temperatures. *BMC Microbiol* **9**: 250.
- Rodríguez-Peña, J.M., García, R., Nombela, C., and Arroyo, J. (2010) The high-osmolarity glycerol (HOG) and cell wall integrity (CWI) signalling pathways interplay: a yeast dialogue between MAPK routes. *Yeast* **27**: 495–502.
- Roels, J.A. (1983) *Energetics and Kinetics in Biotechnology*. Amsterdam, the Netherlands: Elsevier University Press.
- Rohlf, M., Albert, M., Keller, N.P., and Kempken, F. (2007) Secondary chemicals protect mould from fungivory. *Biol Lett* **3**: 523–525.
- Röling, W.F.M., Milner, M.G., Jones, D.M., Fratepietro, F., Swannell, R.P.J., and Head, I.M. (2004) Bacterial community dynamics and hydrocarbon degradation during a field-scale evaluation of bioremediation on a mudflat beach contaminated with buried oil. *Appl Environ Microbiol* **70**: 2603–2613.
- Rosa, C.A., Novak, F.R., Almeida, J.A.G., Mendonça-Hagler, L.C., and Hagler, A.N. (1990) Yeasts from human milk collected in Rio de Janeiro, Brazil. *Rev Microbiol* **21**: 361–363.
- Rose, A.H. (1982) *Fermented Foods*. New York, USA: Academic Press.
- Sadeghi, A.M., Starr, J.L., Teasdale, J.R., Rosecrance, R.C., and Rowland, R.A. (2007) Real-time soil profile water content as influenced by weed-corn competition. *Soil Sci* **172**: 759–769.
- Salisbury, E. (1961) *Weeds and Aliens*. London, UK: Collins.
- Salonen, K., and Rosenberg, M. (2000) Advantages from diel vertical migration can explain the dominance of *Gonyostomum semen* (Raphidophyceae) in a small, steeply-stratified humic lake. *J Plankton Res* **22**: 1841–1853.
- Salvadó, Z., Arroyo-López, F.N., Guillamón, J.M., Salazar, G., Querol, A., and Barrio, E. (2011) Temperature adaptation markedly determines evolution within the genus *Saccharomyces*. *Appl Environ Microbiol* **77**: 2292–2302.
- dos Santos Sant'Ana, G., Paes, L., Vieira Paiva, A.F., Fietto, L.G., Totola, A.H., Trópia, M.J.M., et al. (2009) Protective effect of ions against cell death induced by acid stress in *Saccharomyces*. *FEMS Yeast Res* **9**: 701–712.
- Santos, P.M., Benndorf, D., and Sá-Correia, I. (2004) Insights into *Pseudomonas putida* KT2440 response to

- phenol- induced stress by quantitative proteomics. *Proteomics* **4**: 2640–2652.
- Saponetti, M.S., Bobba, F., Salerno, G., Scarfato, A., Corcelli, A., and Cucolo, A. (2011) Morphological and structural aspects of the extremely halophilic archaeon *Haloquadratum walsbyi*. *PLoS ONE* **6**: e18653.
- Saum, S.H., Pfeiffer, F., Palm, P., Rampp, M., Schuster, S.C., Müller, V., et al. (2012) Chloride and organic osmolytes: a hybrid strategy to cope with elevated salinities by the moderately halophilic, chloride-dependent bacterium *Halobacillus halophilus*. *Environ Microbiol*. doi: 10.1111/j.1462-2920.2012.02770.x.
- Schnitzler, A., and Bailey, J. (2008) Genetic polymorphism and phenotypic plasticity: two advantages for the dispersal of Japanese knotweeds. *Rev Ecol* **63**: 209–217.
- Segura, A., Godoy, P., van Dillewijn, P., Hurtado, A., Arroyo, A., Santacruz, S., and Ramos, J.L. (2005) Proteomic analysis reveals the participation of energy- and stress-related proteins in the response of *Pseudomonas putida* DOT-T1E to toluene. *J Bacteriol* **187**: 5937–5945.
- Selezska, K., Kazmierczak, M., Müsken, M., Garbe, J., Schobert, M., Häussler, S., et al. (2012) *Pseudomonas aeruginosa* population structure revisited under environmental focus: impact of water quality and phage pressure. *Environ Microbiol* **14**: 1952–1967.
- Senthikumar, K., Udaiyan, K., and Manian, S. (1993) Successional pattern of mycoflora associated with litter degradation in a *Cymbopogon caesius*-dominated tropical grassland. *Trop Grasslands* **27**: 121–127.
- Seward, R., Willetts, J.C., Dinsdale, M.G., and Lloyd, D. (1996) The effects of ethanol, hexan-1-ol, and 2-phenylethanol on cider yeast growth, viability, and energy status. *J Inst Brew* **102**: 439–443.
- Shi, X.L., Yang, L.Y., Niu, X.J., Xiao, L., Kong, Z.M., Qin, B.Q., et al. (2003) Intracellular phosphorus metabolism of *Microcystis aeruginosa* under various redox potential in darkness. *Microbiol Res* **158**: 345–352.
- Shukurov, R.R., Voblikova, V.D., Nikonorova, A.K., Komakhin, R.A., Komakhina, V.V., Egorov, T.A., et al. (2012) Transformation of tobacco and Arabidopsis plants with *Stellaria media* genes encoding novel hevein-like peptides increases their resistance to fungal pathogens. *Trans Res* **21**: 313–325.
- Sicard, D., and Legras, J.-L. (2011) Bread, beer and wine: yeast domestication in the *Saccharomyces sensu stricto* complex. *C R Biol* **334**: 229–236.
- Siegel, S.M., and Speitel, T.W. (1976) Performance of fungi in low temperature and hypersaline environments. *Life Sci Space Res* **14**: 351–354.
- Sikkema, J., de Bont, J.A.M., and Poolman, B. (1995) Mechanisms of membrane toxicity of hydrocarbons. *Microbiol Rev* **59**: 201–222.
- Simonetta, M.P., Verga, R., Fretta, A., and Hanozet, G.M. (1989) Induction of D-amino-acid oxidase by D-alanine in *Rhodotorula gracilis* grown in defined medium. *J Gen Microbiol* **135**: 593–600.
- Sinha, M., Srinivasa, H., and Macaden, R. (2007) Antibiotic resistance profile 0026; extended spectrum β -lactamase (ESBL) production in *Acinetobacter* species. *Indian J Med Res* **126**: 63–67.
- Snow, D. (1949) The germination of mould spores at controlled humidities. *Ann Appl Biol* **36**: 1–13.
- Steinkraus, K.H. (ed.) (1983) *Handbook of Indigenous Fermented Foods*. New York, USA: Marcel Dekker.
- Stinson, M.W., and Hayden, C. (1979) Secretion of phospholipase-c by *Pseudomonas aeruginosa*. *Infect Immun* **25**: 558–564.
- Stoyanova-Bakalova, E., and Toncheva-Panova, T. (2003) Subcellular adaptation to salinity and irradiance in *Dunaliella salina*. *Biol Plant* **47**: 233–236.
- Stratford, M. (2006) Food and beverage spoilage yeasts. In *The Yeast Handbook*. Querol, A., and Fleet, G.H. (eds). Berlin, Heidelberg: Springer-Verlag, pp. 335–379.
- Straus, E., and Wu, N. (1980) Radioimmunoassay of tuberculo-protein derived from *Mycobacterium tuberculosis*. *Proc Natl Acad Sci USA* **77**: 4301–4304.
- Suzuki, C., Ando, Y., and Machida, S. (2001) Interaction of SMKT, a killer toxin produced by *Pichia farinosa*, with the yeast cell membranes. *Yeast* **18**: 1471–1478.
- Syn, C.K.C., Magnuson, J.K., Kingsley, M.T., and Swarup, S. (2004) Characterization of *Pseudomonas putida* genes responsive to nutrient limitation. *Microbiology* **150**: 1661–1669.
- Szopinska, A., and Morsomme, P. (2010) Quantitative proteomic approaches and their application in the study of yeast responses. *OMICS* **14**: 639–649.
- Tamang, J.P. (2010) *Himalayan Fermented Foods: Microbiology, Nutrition, and Ethnic Values*. Boca Raton, USA: CRC Press.
- Tark, M., Tover, A., Tarassova, K., Tegova, R., Kivi, G., Hörak, R., et al. (2005) A DNA polymerase V homologue encoded by TOL plasmid pWW0 confers evolutionary fitness on *Pseudomonas putida* under conditions of environmental stress. *J Bacteriol* **187**: 5203–5213.
- Therkildsen, M.S., Isaksen, M.F., and Lomstein, B.A. (1997) Urea production by the marine bacteria *Delvay venusta* and *Pseudomonas stutzeri* grown in a minimal medium. *Aquat Microb Ecol* **13**: 213–217.
- Thomas, K.C., Hynes, S.H., and Ingledew, W.M. (1993) Excretion of proline by *Saccharomyces cerevisiae* during fermentation of arginine-supplemented high gravity wheat mash. *J Ind Microbiol* **12**: 93–98.
- Thomas, R., Grimsley, N., Escande, M.L., Subirana, L., Derelle, E., and Moreau, H. (2011) Acquisition and maintenance of resistance to viruses in eukaryotic phytoplankton populations. *Environ Microbiol* **13**: 1412–1420.
- Thurston, M.I., Pallett, D.W., Cortina-Borja, M., Edwards, M.L., Raybould, A.F., and Cooper, J.I. (2001) The incidence of viruses in wild *Brassica nigra* in Dorset (UK). *Ann Appl Biol* **139**: 277–284.
- Timberlake, W.E., Frizzell, M.A., Richards, K.D., and Gardner, R.C. (2011) A new yeast genetic resource for analysis and breeding. *Yeast* **28**: 63–80.
- Timmis, K.N. (2002) *Pseudomonas putida*: a cosmopolitan opportunist par excellence. *Environ Microbiol* **4**: 779–781.
- Timmis, K.N. (ed.) (2009) *Handbook of Hydrocarbon and Lipid Microbiology – Hydrocarbons, Oils and Lipids: Diversity, Properties and Formation*, Vol. 2. New York, USA: Springer.

- Tiwari, B.K., Valdramidis, V.P., O'Donnell, C.P., Muthukumarappan, K., Bourke, P., and Cullen, P.J. (2009) Application of natural antimicrobials for food preservation. *J Agric Food Chem* **57**: 5987–6000.
- Tobin, M.B., Peery, R.B., and Skatrud, P.L. (1997) Genes encoding multiple drug resistance-like proteins in *Aspergillus fumigatus* and *Aspergillus flavus*. *Gene* **200**: 11–23.
- Tobin, K.M., McGrath, J.W., Mullan, A., Quinn, J.P., and O'Connor, K.E. (2007) Polyphosphate accumulation by *Pseudomonas putida* CA-3 and other medium-chain-length polyhydroxyalkanoate-accumulating bacteria under aerobic growth conditions. *Appl Environ Microbiol* **73**: 1383–1387.
- Todd, P.P., Lucero, C.A., and Falkinham, J.O. (2004) Health impacts of environmental mycobacteria. *Clin Microbiol Rev* **17**: 98–106.
- Toh, H., Kayingo, G., van der Merwe, M.J., Kilian, S.G., Hallsworth, J.E., Hohmann, S., et al. (2001) Implications of *FPS1* deletion and membrane ergosterol content for glycerol efflux from *Saccharomyces cerevisiae*. *FEMS Yeast Res* **1**: 205–211.
- Trigal, C., Goedkoop, W., and Johnson, R.K. (2011) Changes in phytoplankton, benthic invertebrate and fish assemblages of boreal lakes following invasion by *Gonyostomum semen*. *Freshw Biol* **56**: 1937–1948.
- Trindade, R.C., Resende, M.A., Silva, C.M., and Rosa, C.A. (2002) Yeasts associated with fresh and frozen pulps of Brazilian tropical fruits. *Syst Appl Microbiol* **25**: 294–300.
- Tsirogiani, E., Aivaliotis, M., Papasotiriou, D.G., Karas, M., and Tsiotis, G. (2006) Identification of inducible protein complexes in the phenol degrader *Pseudomonas* sp. strain phDV1 by blue native gel electrophoresis and mass spectrometry. *Amino Acids* **30**: 63–72.
- Tsubo, M., Nishihara, E., Nakamatsu, K., Cheng, Y.X., and Shinoda, M. (2012) Plant volatiles inhibit restoration of plant species communities in dry grassland. *Basic Appl Ecol* **13**: 76–84.
- Turk, M., Plemenitaš, A., and Gunde-Cimerman, N. (2011) Extremophilic yeasts: plasma-membrane fluidity as determinant of stress tolerance. *Fungal Biol* **115**: 950–958.
- Vallentyne, J.R. (1963) Environmental biophysics and microbial ubiquity. *Ann N Y Acad Sci* **108**: 342–352.
- Vasileva-Tonkova, E., and Gesheva, V. (2007) Biosurfactant production by Antarctic facultative anaerobe *Pantoea* sp. during growth on hydrocarbons. *Curr Microbiol* **54**: 136–141.
- Visser, A.A., Kooij, P.W., Debets, A.J.M., Kuyper, T.W., and Aanen, D.K. (2011) *Pseudoxylaria* as stowaway of the fungus-growing termite nest: interaction asymmetry between *Pseudoxylaria*, *Termitomyces* and free-living relatives. *Fungal Ecol* **4**: 322–332.
- Vital, M.J.S., Abranches, J., Hagler, A.N., and Mendonça-Hagler, L.C. (2002) Mycocinogenic yeasts isolated from Amazon soils of the Maracá Ecological Station, Roraima-Brazil. *Braz J Microbiol* **33**: 230–235.
- Volkers, R.J.M., de Jong, A.L., Hulst, A.G., van Baar, B.L.M., de Bont, J.A.M., and Wery, J. (2006) Chemostat-based proteomic analysis of toluene-affected *Pseudomonas putida* S12. *Environ Microbiol* **8**: 1674–1679.
- Volkers, R.J.M., Ballerstedt, H., Ruijsenaars, H.J., de Bont, J.A., de Winde, J.H., and Wery, J. (2009) *Trgl*, toluene repressed gene I, a novel gene involved in toluene-tolerance in *Pseudomonas putida* S12. *Extremophiles* **13**: 283–297.
- Volkers, R.J.M., Ballerstedt, H., de Winde, J.H., and Ruijsenaars, H.J. (2010) Isolation and genetic characterization of an improved benzene-tolerant mutant of *Pseudomonas putida* S12. *Environ Microbiol Rep* **2**: 456–460.
- Walker, G.M. (2011) *Pichia anomala*: cell physiology and biotechnology relative to other yeasts. *Antonie Van Leeuwenhoek* **99**: 25–34.
- Wang, R.-L., Staehelin, C., Peng, S.-L., Wang, W.-T., Xie, X.-M., and Lu, H.-N. (2010) Responses of *Mikania micrantha*, an invasive weed to elevated CO₂: induction of β -caryophyllene synthase, changes in emission capability and allelopathic potential of β -caryophyllene. *J Chem Ecol* **36**: 1076–1082.
- Ward, O.P., Qin, W.M., Dhanjoon, J., Ye, J., and Singh, A. (2006) Physiology and biotechnology of *Aspergillus*. *Adv Appl Microbiol* **58**: 1–75.
- Watanabe, A., Yano, K., Ikebukuro, K., and Karube, I. (1998) Cyanide hydrolysis in a cyanide-degrading bacterium, *Pseudomonas stutzeri* AK61, by cyanidase. *Microbiology* **144**: 1677–1682.
- Wecke, T., Bauer, T., Harth, H., Mäder, U., and Mascher, T. (2011) The rhamnolipid stress response of *Bacillus subtilis*. *FEMS Microbiol Lett* **323**: 113–123.
- Wendell, D.L., and Bisson, L.F. (1994) Expression of high-affinity glucose transport protein Hxt2p of *Saccharomyces cerevisiae* is both repressed and induced by glucose and appears to be regulated post-translationally. *J Bacteriol* **176**: 3730–3737.
- Westenberg, B., Boller, T., and Wiemken, A. (1989) Lack of arginine- and polyphosphate-storage pools in a vacuole-deficient mutant (endl) of *Saccharomyces cerevisiae*. *FEBS Lett* **254**: 133–136.
- Wheeler, K.A., and Hocking, A.D. (1993) Interactions among xerophilic fungi associated with dried salted fish. *J Appl Bacteriol* **74**: 164–169.
- Wiesenborn, D.P., Rudolph, F.B., and Papoutsakis, E.T. (1989) Phosphotransbutyrylase from *Clostridium acetobutylicum* ATCC-824 and its role in acidogenesis. *Appl Environ Microbiol* **55**: 317–322.
- Williams, J.P., and Hallsworth, J.E. (2009) Limits of life in hostile environments: no barriers to biosphere function? *Environ Microbiol* **11**: 3292–3308.
- de Wit, R., and Bouvier, T. (2006) 'Everything is everywhere, but, the environment selects'; what did Baas Becking and Beijerinck really say? *Environ Microbiol* **8**: 755–758.
- Wu, S.J., Zhang, H.X., Yu, X., and Chen, J.M. (2009) Identification and cloning of a gene encoding dichloromethane dehalogenase from a methylotrophic bacterium, *Bacillus circulans* WZ-12 CCTCC M 207006. *Bioprocess Biosyst Eng* **32**: 845–852.
- Wu, T.-H., Liao, M.-H., Kuo, W.-Y., Huang, C.-H., Hsieh, H.-L., and Jinn, T.-L. (2011) Characterization of copper/zinc and manganese superoxide dismutase in green bamboo (*Bambusa oldhamii*): cloning, expression and regulation. *Plant Physiol Biochem* **49**: 195–200.

- Xing, W., Huang, W.-M., Li, D.-H., and Liu, Y.-D. (2007) Effects of iron on growth, pigment content, photosystem II efficiency, and siderophores production of *Microcystis aeruginosa* and *Microcystis wesenbergii*. *Curr Microbiol* **55**: 94–98.
- Xuan, T.D., Chung, M.I., Khanh, T.D., and Tawata, S. (2006) Identification of phytotoxic substances from early growth of barnyard grass (*Echinochloa crusgalli*) root exudates. *J Chem Ecol* **32**: 895–906.
- Yalcin, S.K., and Ozbas, Z.Y. (2008) Effects of pH and temperature on growth and glycerol production kinetics of two indigenous wine strains of *Saccharomyces cerevisiae* from Turkey. *Braz J Microbiol* **39**: 325–332.
- Yancey, P.H. (2005) Organic osmolytes as compatible, metabolic and counteracting cytoprotectants in high osmolarity and other stresses. *J Exp Biol* **208**: 2819–2830.
- Yang, Z., and Kong, F. (2012) Formation of large colonies: a defense mechanism of *Microcystis aeruginosa* under continuous grazing pressure by flagellate *Ochromonas* sp. *J Limnol* **71**: 61–66.
- Zhu, H.-Y., Xu, H., Dai, X.-Y., Zhang, Y., Ying, H.-J., and Ouyang, P.-K. (2010) Production of D-arabitol by a newly isolated *Kodamaea ohmeri*. *Bioproc Biosyst Eng* **33**: 565–571.
- Zuroff, T.R., and Curtis, W.R. (2012) Developing symbiotic consortia for lignocellulosic biofuel production. *Appl Microbiol Biotechnol* **93**: 1423–1435.

Supporting information

Additional Supporting Information may be found in the online version of this article:

Table S1. Examples of open habitats in which specific microbes attain dominance.

Table S2. Biology of microbial weeds compared with that of other ecophysiological groups.